

# Circular (Yet Sound) Proofs in Propositional Logic

Albert Atserias  
Universitat Politècnica de Catalunya

Massimo Lauria  
Sapienza - Università di Roma

April 11, 2023

## Abstract

Proofs in propositional logic are typically presented as trees of derived formulas or, alternatively, as directed acyclic graphs of derived formulas. This distinction between tree-like vs. dag-like structure is particularly relevant when making quantitative considerations regarding, for example, proof size. Here we analyze a more general type of structural restriction for proofs in rule-based proof systems. In this definition, proofs are directed graphs of derived formulas in which cycles are allowed as long as every formula is derived at least as many times as it is required as a premise. We call such proofs “circular”. We show that, for all sets of standard inference rules with single or multiple conclusions, circular proofs are sound. We start the study of the proof complexity of circular proofs at Circular Resolution, the circular version of Resolution. We immediately see that Circular Resolution is stronger than Dag-like Resolution since, as we show, the propositional encoding of the pigeonhole principle has circular Resolution proofs of polynomial size. Furthermore, for derivations of clauses from clauses, we show that Circular Resolution is, surprisingly, equivalent to Sherali-Adams, a proof system for reasoning through polynomial inequalities that has linear programming at its base. As corollaries we get: 1) polynomial-time (LP-based) algorithms that find Circular Resolution proofs of constant width, 2) examples that separate Circular from Dag-like Resolution, such as the pigeonhole principle and its variants, and 3) exponentially hard cases for Circular Resolution. Contrary to the case of Circular Resolution, for Frege we show that circular proofs can be converted into tree-like proofs with at most polynomial overhead.

## 1 Introduction

Logical proofs are traditionally presented as sequences of formulas where each formula is either a hypothesis, or is deduced from some previous formulas in the sequence by the means of some inference step. In rule-based proofs systems each inference step is achieved by instantiating one in some specific and finite set of inference rules. Equivalently, any such proof can be represented by a directed acyclic graph, or *dag*, with one vertex for each formula in the sequence, and edges pointing forward from the premises to the conclusions of each inference rule application.

In this paper we discuss an alternative and more general way of composing proofs: we allow cycles in the graph. In general, and not suprisingly, unlimited circular reasoning of this type may be unsound. However,

---

A preliminary shorter version of this paper was published in the Proceedings of 22nd International Conference on Theory and Applications of Satisfiability Testing (SAT 2019), Lecture Notes in Computer Science 11628, Springer 2019, ISBN 978-3-030-24257-2, pp. 1-18, Lisbon, Portugal, July 9-12, 2019. The full version of this paper has been published in ACM Trans. Comput. Logic. 24(3):1–20. This pre-print version can also be found at CoRR abs/1802.05266.

when every formula is derived at least as many times as it is used as a premise of an inference step, we show that soundness is guaranteed. Hence we appropriately call these objects with the name of *circular proofs*.

More formally, our soundness requirement is phrased in terms of *flow assignments*: each rule application must carry a *flow*, a positive integer which intuitively means that in order to produce that many copies of each conclusion of the rule we must have produced at least that many copies of each premise. Flow assignments induce a notion of *balance* of a formula in the proof, which is the difference between the number of times that the formula is produced as a conclusion and the number of times that it is required as a premise. Given these definitions, a proof-graph will be an actual circular proof if it admits a flow assignment that satisfies the following *flow-balance* condition: the only formulas of strictly negative balance are the hypotheses, and the goal formulas display strictly positive balance. With this interpretation of flows, circular proofs have the appealing flavour of a network in which demands are fulfilled by the hypotheses, and flow towards the goal formulas, which produce surplus. Accordingly, and in analogy with the theory of classical network flows [29], it makes no difference whether the flows are required to be integers or real numbers, and valid flow assignments can be found efficiently, when they exist, by linear programming techniques.

While proof-graphs with unrestricted cycles are, in general, unsound, we show that circular proofs *are* sound. We prove this in two ways. The first one is combinatorial in nature and is phrased in the style of traditional soundness proofs in standard proof systems. Concretely, given a truth assignment that falsifies a goal formula, the soundness proof constructs a path of falsified formulas until it reaches a hypothesis, and does so by induction on the total flow-sum of the flow assignment that satisfies the flow-balance condition. This proof is more informative and intuitive, but it is also inefficient in the sense that the process of building the path is not polynomial in the size of the proof. The second proof is (semi-)algebraic and is phrased in the style of the duality theorem for linear programming. Concretely, we phrase the unsoundness of the proof as the feasibility of a linear program and observe that the existence of a flow assignment that satisfies the flow-balance condition gives a witness of its infeasibility. This second proof is less intuitive but can be efficiently simulated by non-circular argument in relatively strong proof systems. It will be useful when trying to understand the power of circular reasoning.

**Proof complexity of circular proofs** With all the definitions in place, we proceed to the study of the power of circular proofs from the perspective of propositional proof complexity.

For Resolution, we show that circularity *does* make a real difference. First we show that the standard propositional formulation of the pigeonhole principle has Circular Resolution proofs of polynomial size. This is in sharp contrast with the well-known fact that Resolution *cannot count*, and that the pigeonhole principle is exponentially hard for (tree-like and dag-like) Resolution [21]. Second we observe that the LP-based proof of soundness of Circular Resolution can be formalized in the Sherali-Adams proof system (with twin variables), which is a proof system for reasoning with polynomial inequalities that has linear programming at its base [31]. Sherali-Adams was originally conceived as a hierarchy of linear programming relaxations for integer programs, but it has also been studied from the perspective of proof complexity in recent years [14, 15, 27, 3].

Surprisingly, it turns out that the converse simulation is also true! For deriving clauses from clauses, Sherali-Adams proofs translate efficiently into Circular Resolution proofs. Moreover, both translations, the one from Circular Resolution into Sherali-Adams and its converse, are efficient in terms of their natural parameters: length/size and width/degree. As corollaries we obtain for Circular Resolution all the proof complexity-theoretic properties that are known to hold for Sherali-Adams: 1) a polynomial-time (LP-based) proof search

algorithm for proofs of bounded width, 2) length-width relationships, 3) separations from dag-like length and width, and 4) explicit exponentially hard examples.

Going beyond resolution we address the question of how circularity affects more powerful propositional proof systems. For Frege systems, which operate with arbitrary propositional formulas through the standard textbook inference rules, we show that circularity adds no power: the circular, dag-like and tree-like variants of Frege polynomially simulate one another. The equivalence between the dag-like and tree-like variants of Frege is well-known [23]; here we add the circular variant to the list. We prove this by formalizing the LP-based proof of soundness for Circular Frege within Tree-like Frege itself. To achieve this we make strong use of the formalization of linear arithmetic in Frege that was developed by Buss in order to get efficient Frege proofs of the pigeonhole principle [12], and that was developed further by Goerdt to show that Tree-like Frege simulates the Cutting Planes proof system [19].

**Earlier work** The idea of allowing cyclic, circular or non-wellfounded proofs has been studied by several communities since at least 20 years ago, from modal  $\mu$ -calculus [26], to predicate logic with inductive definitions [10, 11], fragments of arithmetic [33, 16], provability logics [30], and linear logic [18]. For classical propositional logic proper and in the context of proof complexity, we are not aware of any work on cyclic, circular, or non-wellfounded proofs that appeared earlier than the conference version of this paper [4]. It seems that our flow-based definition of circular proofs had not been considered before.

Niwiński and Walukiewicz [26] introduced an infinitary tableau method for the modal  $\mu$ -calculus. The proofs are regular infinite trees that are represented by finite graphs with cycles, along with a decidable *progress condition* on the cycles to guarantee their soundness. A sequent calculus version of this tableau method was proposed in [17], and explored further in [34]. In his PhD thesis, Brotherston [10] introduced a *cyclic* proof system for the extension of first-order logic with inductive definitions; see also [11] for a journal article presentation of the results. The proofs in [11] are ordinary proofs of the first-order sequent calculus extended with the rules that define the inductive predicates, along with a set of *backedges* that link equal formulas in the proof. The soundness is guaranteed by an additional *infinite descent condition* along the cycles that is very much inspired by the progress condition in Niwiński-Walukiewicz’ tableau method. We refer the reader to Section 8 from [11] for a careful overview of the various flavours of proofs with cycles for logics with inductive definitions.

Shoemith and Smiley [32] initiate the study of inference based propositional proofs with multiple conclusions. In order to do so they introduce a graphical representation of proofs where nodes represent either formulas or inference steps, in a way similar to our definition in Section 2. While most of that book does not consider proof with cycles, in Section 10.5 they do mention briefly this possibility but they do not analyze it any further.

The Sherali-Adams hierarchy of linear programming relaxations has received considerable attention in recent years for its relevance to combinatorial optimization and approximation algorithms; see the original [31], and [5] for a recent survey. In its original presentation, the Sherali-Adams hierarchy can already be thought of as a proof system for reasoning with polynomial inequalities, with the levels of the hierarchy corresponding to the degrees of the polynomials. For propositional logic, the system was studied in [14], and developed further in [27, 3]. Those works consider the version of the proof system in which each propositional variable  $X$  comes with a formal *twin variable*  $\bar{X}$ , that is to be interpreted by the negation of  $X$ . This is the version of Sherali-Adams that we use. It was already known from [15] that this version of the Sherali-Adams proof system polynomially simulates standard Resolution, and has polynomial-size proofs of the pigeonhole principle.

## 2 Preliminaries

### 2.1 Formulas

A *literal* is a variable  $X$  or the negation of a variable  $\bar{X}$ ; we also say that literal  $\bar{X}$  is the negation of literal  $X$ , and vice-versa. The class of formulas in *negation normal form* is the smallest class of formulas that contains the literals and is closed under conjunction  $\wedge$  and disjunction  $\vee$ . If  $A$  is a formula in negation normal form, we write  $\bar{A}$  for its dual formula, which is defined recursively as follows: If  $A$  is a literal, then  $\bar{A}$  is its negation. If  $A = B \vee C$ , then  $\bar{A} = \bar{B} \wedge \bar{C}$ . If  $A = B \wedge C$ , then  $\bar{A} = \bar{B} \vee \bar{C}$ . Note that the dual of the dual of  $A$  is  $A$  itself. A *truth-assignment* is a mapping that assigns a truth-value *true* (1) or *false* (0) to each variable. Truth-assignments evaluate formulas in the natural way through the standard interpretations of negation, conjunction, and disjunction. The *empty formula* is denoted by 0, and is always false by convention. Its complement  $\bar{0}$  is denoted by 1 and is always true by convention. If a truth-assignment evaluates a formula to true, then we say that it satisfies it. A *substitution* is a mapping that assigns a formula to each variable. Applying a substitution to a formula means replacing all variables by the formulas to which they are mapped to by the substitution, simultaneously all at once.

We think of disjunction as binding *sets* of formulas, or, equivalently, as a binary operation on formulas that is associative, commutative and idempotent. This means that the formula  $(A \vee B) \vee C$  is considered the same as  $A \vee (B \vee C)$ , which we just write as  $A \vee B \vee C$ . Also the formula  $A \vee B$  is considered the same as  $B \vee A$ , and the formula  $A \vee A$  is considered the same as  $A$ . Similarly, we view conjunction as a binary operation on formulas that is associative, commutative and idempotent. The empty formula 0 and its complement 1 are the neutral elements of  $\vee$  and  $\wedge$ , respectively. Thus the formulas  $0 \vee A$  and  $1 \wedge A$  are considered the same as  $A$ . These conventions about disjunctions and conjunctions mean that our syntax for formulas in negation normal form could have been defined as follows: (1) every literal is a formula, (2) if  $S$  is a set of formulas none of which starts with  $\vee$ , then  $\vee S$  is a formula, (3) if  $S$  is a set of formulas none of which starts with  $\wedge$ , then  $\wedge S$  is a formula, and (4) nothing else is a formula. The empty formula 0 and its complement 1 are taken to be  $\vee \emptyset$  and  $\wedge \emptyset$ , respectively. We adopt this *unbounded fan-in* definition of syntax, but continue to use the notation  $A_1 \vee \dots \vee A_n$  even if the  $A_i$  may be disjunctions themselves. The *size*  $s(A)$  of a formula  $A$  is defined inductively: if  $A$  is a literal, then  $s(A) = 1$ , and if  $A = \vee S$  or  $A = \wedge S$ , then  $s(A) = 1 + \sum_{B \in S} s(B)$ .

An *elementary tautology* is a formula of the form  $\bar{A} \vee A$ , where  $A$  is a formula. Note that by the definition of the dual of a formula (and the convention to read disjunctions up to associativity), the formula  $\bar{A} \vee \bar{B} \vee (A \wedge B)$  is an elementary tautology. If  $\Gamma$  is a set of formulas, a disjunction of formulas in  $\Gamma$  is a formula of the form  $A_1 \vee \dots \vee A_m$ , where  $m$  is a non-negative integer and each  $A_i$  is a formula in  $\Gamma$ . Disjunctions of formulas in  $\Gamma$  are also called  $\Gamma$ -clauses or  $\Gamma$ -cedents. A clause is a disjunction of literals.

### 2.2 Inference-Based Proofs

An *inference rule* is given by a sequence of *premise formulas*  $A_1, \dots, A_r$  and a sequence of *conclusion formulas*  $B_1, \dots, B_s$  with the property that every truth assignment that satisfies all the premises also satisfies all the conclusions. Here are four important examples:

$$\frac{}{A \vee \bar{A}} \quad \frac{C \vee A \quad D \vee \bar{A}}{C \vee D} \quad \frac{C \vee A \quad D \vee B}{C \vee D \vee (A \wedge B)} \quad \frac{C}{C \vee D}. \quad (1)$$

These are the standard inference rules of a Tait-style calculus for propositional logic [35]. The rules are called *axiom*, *cut*, *introduction of conjunction*, and *weakening*, respectively. An *instance* of an inference rule




Figure 1: The directed acyclic graph representation of a proof of  $A_7$  from the set of hypothesis formulas  $A_1$  and  $A_2$  through the inference rules  $R_1, \dots, R_5$ . Formula-vertices are represented by boxes and inference-vertices are represented by circles. Formula  $A_3$  is used twice as the premise of an inference, and  $A_5$  is produced twice as the conclusion of an inference. All rules except  $R_4$  have exactly one conclusion formula;  $R_4$  has two. All rules except  $R_2$  have at least one premise formula;  $R_2$  has none.

is obtained from applying a substitution to its variables. Note that every instance of a rule is a rule itself, which has its own premise formulas and conclusion formulas.

In almost all classical examples in the literature, inference rules have a single conclusion formula. The reason for this is that for classical (i.e., non-circular) proofs one may simply split a rule with  $s$  conclusion formulas into  $s$  different single-conclusion rules, with little conceptual change. However, for circular proofs a specific rule with two conclusion formulas will play an important role; this is the *symmetric weakening*, or *split*, rule:

$$\frac{C}{C \vee A \quad C \vee \bar{A}}. \quad (2)$$

When we apply (2) we say that we split  $C$  on  $A$ . In all these examples the formulas  $C$ ,  $D$ , and  $A$  could be a single literal, the empty formula 0, or its complement 1.

Fix a set  $\mathcal{R}$  of inference rules, a set  $A_1, \dots, A_m$  of *hypothesis formulas*, and a *goal formula*  $A$ . A *proof of  $A$  from  $A_1, \dots, A_m$* , also called a *derivation*, is a finite sequence of formulas that ends in  $A$  and such that each formula in the sequence is either contained in  $A_1, \dots, A_m$ , or is one of the conclusion formulas of an instance of an inference rule in  $\mathcal{R}$  that has all its premise formulas appearing earlier in the sequence. A derivation of  $A$  from *nothing* is also called a *proof of  $A$* . A *refutation of  $A_1, \dots, A_m$*  is a derivation of the empty formula 0 from  $A_1, \dots, A_m$ . The *length* of the derivation is the length of the sequence, and its *size* is the sum of the sizes of the formulas in the sequence.

Proofs and derivations are naturally represented through directed acyclic graphs, a.k.a. *dags*; see Figure 1. The graph has one *formula-vertex* for each formula in the sequence, and one *inference-vertex* for each inference step that produces a formula in the sequence. Each formula-vertex is labelled by the corresponding formula, and each inference-vertex is labelled by the corresponding instance of the corresponding inference rule. Each inference-vertex that is labelled by an inference rule that has  $r$  premise formulas and  $s$  conclusion formulas has, accordingly,  $r$  incoming edges from the corresponding premise formula-vertices, and at least one and at most  $s$  outgoing edges towards the corresponding conclusion formula-vertices. The directed acyclic graph of a proof  $\Pi$  is its *proof-graph*, and is denoted by  $G(\Pi)$ . A proof  $\Pi$  is called *tree-like* if  $G(\Pi)$  is a tree.

### 2.3 Frege and Resolution Proof Systems

An inference-based proof system is given by a set of allowed inference rules, a set of allowed formulas, and a set of allowed proof-graphs. Two typical sets of allowed proof-graphs are the set of dags, for *dag-like* proofs, and the set of trees, for *tree-like* proofs. If the set of allowed proof-graphs is omitted, dag-like is assumed by default. A proof system  $P$  is said to *polynomial simulate* another proof system  $P'$  if there is a polynomial-time algorithm that, given a proof  $\Pi'$  in  $P'$  as input, computes a proof  $\Pi$  in  $P$ , such that  $\Pi$  has the same goal formula and the same hypothesis formulas as  $\Pi'$ . *Frege* and *Resolution* are both inference-based proof systems, as defined next.

In our definition of Frege the set of allowed inference rules are axiom, cut, introduction of conjunction, and weakening as defined in (1), and the set of allowed formulas is the set of all formulas in negation normal form. Being equivalent to a Tait-style calculus, our definition of Frege is sound and (implicationally) complete for formulas in negation normal form. This means that if  $A$  has a Frege proof from the set of hypothesis formulas  $A_1, \dots, A_m$ , then every truth assignment that satisfies all the formulas in  $A_1, \dots, A_m$  also satisfies  $A$ , and vice-versa.

In our definition of Resolution the only allowed inference rule is cut and the allowed formulas are the clauses. This proof system is sound and complete as a refutation system. This means that if the set of clauses  $A_1, \dots, A_m$  has a Resolution refutation, then there is no truth-assignment that satisfies all clauses  $A_1, \dots, A_m$  simultaneously, and vice-versa. In order to turn Resolution into a proof system that is sound and complete for deriving clauses from clauses, one needs to add the axiom and weakening rules to the set of allowed rules. The *width* of a Resolution proof is the number of literals of its largest clause.

### 2.4 Frege and Resolution with Symmetric Rules

Consider an inference-based proof system in which elementary tautologies of the form  $A \vee \bar{A}$  may be introduced at any point in the proof through the axiom rule, and that in addition has the following two nicely symmetric-looking inference rules:

$$\frac{C \vee A \quad C \vee \bar{A}}{C} \qquad \frac{C}{C \vee A \quad C \vee \bar{A}}. \quad (3)$$

These rules are called *symmetric cut* and *symmetric weakening*, or *split*, respectively. Note the subtle difference between the symmetric cut rule and the standard cut rule in (1): in the symmetric cut rule, both premise formulas have the same *side formula*  $C$ . This difference is minor: an application of the non-symmetric cut rule that derives  $C \vee D$  from  $C \vee A$  and  $D \vee \bar{A}$  may be efficiently simulated as follows (here and in what follows, the applicability of the rules has to be read up to associativity, symmetry, and idempotency of disjunctions and conjunctions, and the second conclusion of the split rule has been suppressed from the list of derived formulas whenever it is not needed):

1.  $C \vee A \vee D$       by split on  $C \vee A$ ,
2.  $D \vee \bar{A} \vee C$       by split on  $D \vee \bar{A}$ ,
3.  $C \vee D$               by symmetric cut on 1 and 2.

Note also that the rules in (3) do not include a rule for *introduction of conjunction* as in (1). In the presence of the elementary tautologies (or, equivalently, the axiom rule), this difference is again minor: an application of the introduction of conjunction rule that derives  $C \vee D \vee (A \wedge B)$  from  $C \vee A$  and  $D \vee B$  may be efficiently simulated by the following sequence:

- |    |   |                              |
|----|---|------------------------------|
| 1. | $\bar{A} \vee \bar{B} \vee (A \wedge B)$        | as an elementary tautology,  |
| 2. | $\bar{A} \vee \bar{B} \vee (A \wedge B) \vee C$ | by split on 1,               |
| 3. | $C \vee A \vee \bar{B} \vee (A \wedge B)$       | by split on $C \vee A$ ,     |
| 4. | $C \vee \bar{B} \vee (A \wedge B)$              | by symmetric cut on 2 and 3, |
| 5. | $C \vee \bar{B} \vee (A \wedge B) \vee D$       | by split on 4,               |
| 6. | $D \vee B \vee C \vee (A \wedge B)$             | by split on $D \vee B$ ,     |
| 7. | $C \vee D \vee (A \wedge B)$                    | by symmetric cut on 5 and 6. |

Thus, for all practical purposes, the Frege proof system as defined in the previous section and the proof system defined here are equivalent. The same observation applies to Resolution. In this case the elementary tautologies are of the form  $X \vee \bar{X}$ , where  $X$  is a variable, and the instances of the symmetric cut and split rules in (3) have a variable for its *cut formula*  $A$ . Note that an application of the standard weakening rule that derives the clause  $C \vee D$  from the clause  $C$  may be efficiently simulated by  $|D|$  many applications of the split rule by introducing one literal at a time; here  $|D|$  denotes the number of literals in  $D$ .

## 2.5 Sherali-Adams Proof System

Let  $X_1, \dots, X_n$  be variables that are intended to range over  $\{0, 1\}$ , and let  $\bar{X}_1, \dots, \bar{X}_n$  be their *twins*, with the intended meaning that  $\bar{X}_i = 1 - X_i$ . Let  $A_1, \dots, A_m$  and  $A$  be polynomials on the variables  $X_1, \dots, X_n$  and  $\bar{X}_1, \dots, \bar{X}_n$ . A *Sherali-Adams proof of  $A \geq 0$  from  $A_1 \geq 0, \dots, A_m \geq 0$*  is a polynomial identity of the form

$$\sum_{j=1}^t Q_j P_j = A, \quad (4)$$

where each  $Q_j$  is a non-negative linear combination of monomials on the variables  $X_1, \dots, X_n$  and  $\bar{X}_1, \dots, \bar{X}_n$ , and each  $P_j$  is a polynomial among  $A_1, \dots, A_m$  or one among the following set of *basic* polynomials:

$$X_i - X_i^2, \quad 1 - X_i - \bar{X}_i, \quad X_i^2 - X_i, \quad X_i + \bar{X}_i - 1, \quad 1. \quad (5)$$

Observe that all basic polynomials, as well as all  $Q_j$ 's, being non-negative linear combinations of monomials, are non-negative on  $\{0, 1\}$ . If  $A_1, \dots, A_m$  are also non-negative, then, by (4),  $A$  must be non-negative. It follows that the proof system is sound. By Theorem 3 in the original paper by Sherali and Adams [31], the proof system is complete for deriving linear inequalities from linear inequalities. Therefore, when clauses are encoded by linear inequalities in the natural way, the proof system is also complete (see also Lemma 4.2 in [3] and also Section 4 in this paper). In a Sherali-Adams proof each  $Q_j$  is given explicitly as a positive linear combination of monomials, where the coefficients in the linear combination are rational numbers written in binary. It follows that the identity asserted by equation (4) can be checked in polynomial time with respect to the length of the proof itself. These three facts together imply that Sherali-Adams is a Cook-Reckhow proof system.

The *degree* of the proof is the maximum of the degrees of the polynomials  $Q_j P_j$  in (4). The *monomial size* of the proof is the sum of the monomial sizes of the polynomials  $Q_j P_j$  in (4), where the monomial size of a polynomial is the number of monomials with non-zero coefficient in its unique representation as a linear combination of monomials. The *bit size* of the proof is the sum of the bit sizes of the polynomials  $Q_j P_j$  in (4), where the bit size of a polynomial is the sum of the bit sizes of its terms, where the bit size of a term is the number of bits that it takes to describe the monomial and to write the rational coefficient in binary.




Figure 2: A circular pre-proof. The vertex labelled  $R_1$  has two premises,  $A_1$  and  $B_1$ , and one conclusion,  $B_1$  itself. The vertex labelled  $R_3$  has two premises,  $B_1$  and  $A_2$ , and one conclusion,  $A_3$ . The vertex labelled  $R_2$  has no premises and one conclusion,  $A_2$ .




Figure 3: An unsound circular pre-proof: the false empty formula  $0$  is derived from no hypotheses. We note that if we were to assign positive weights to the inference-vertices, then it would always be the case that the sum of the weights that enter  $X$  minus the sum of the weights that leave  $X$  would always be negative (and the same for  $\bar{X}$ ). As we will see, this turns out to be the *only reason* for it not being sound.

### 3 Circular Proofs

Informally, a circular proof will be defined as a “proof with cycles”. Formally such objects will be called *circular pre-proofs* because, in general, they are not sound. We define *circular proofs* by adding a global yet efficiently checkable requirement on the definition of pre-proof that guarantees its soundness.

#### 3.1 Definition

A *circular pre-proof* is a directed graph with two types of vertices: formula-vertices and inference-vertices. All edges of the graph go from a formula-vertex to an inference-vertex, or from an inference-vertex to a formula-vertex. Thus, the graph is *bipartite*. Each formula-vertex is labelled by a formula, and each inference-vertex is labelled by an instance of an inference rule that has the formulas that label its in-neighbors as premises, and the formulas that label its out-neighbors as conclusions. If  $\Pi$  is a pre-proof, we use  $G(\Pi)$  to denote the underlying bipartite graph, ignoring the labels. When  $\Pi$  is clear from the context, we write  $I$  and  $J$  for the sets of inference- and formula-vertices of  $G(\Pi)$ , respectively, and  $N^-(u)$  and  $N^+(u)$ , respectively, for the sets of in- and out-neighbours of a vertex  $u$  of  $G(\Pi)$ . Figure 2 illustrates these definitions.

By the correspondence between proofs and their dags as defined in Section 2, a circular pre-proof whose underlying graph is a dag is just the same as an ordinary proof. However, general circular pre-proofs need not be sound; see Figure 3 for an example of an unsound circular pre-proof. In order to ensure soundness we need to require a global condition as defined next.

A *flow assignment* for a circular pre-proof  $\Pi$  is an assignment  $F : I \rightarrow \mathbb{R}^+$  of positive real weights, or

flows, where  $I$  is the set of inference-vertices of the graph  $G(\Pi)$  of  $\Pi$ . The flow-extended graph that labels each inference-vertex  $w$  of  $G(\Pi)$  by its flow  $F(w)$  is denoted by  $G(\Pi, F)$ . The *inflow* of a formula-vertex in  $G(\Pi, F)$  is the sum of the flows of its in-neighbours. Similarly, the *outflow* of a formula-vertex in  $G(\Pi, F)$  is the sum of the flows of its out-neighbours. The *balance* of a formula-vertex  $u$  of  $G(\Pi, F)$  is the inflow minus the outflow of  $u$ , and is denoted by  $B(u)$ . In symbols,

$$B(u) := \sum_{w \in N^-(u)} F(w) - \sum_{w \in N^+(u)} F(w). \quad (6)$$

This notion allows us to define *sources* and *sinks* in  $G(\Pi, F)$ . The formula-vertices of strictly negative balance are the sources of  $G(\Pi, F)$ , and those of strictly positive balance are the sinks of  $G(\Pi, F)$ . We think of flow assignments as witnessing a proof of a formula that labels a sink from the set of formulas that label the sources. Concretely, for a given set of hypothesis formulas  $\mathcal{H}$  and a given goal formula  $A$ , we say that the flow assignment *witnesses a proof of  $A$  from  $\mathcal{H}$*  if every source of  $G(\Pi, F)$  is labelled by a formula in  $\mathcal{H}$ , and some sink of  $G(\Pi, F)$  is labelled by the formula  $A$ .

Finally, a *circular proof of  $A$  from  $\mathcal{H}$*  is a circular pre-proof for which there exists a flow assignment that witnesses a proof of  $A$  from  $\mathcal{H}$ . The *length* of a circular proof  $\Pi$  is the number of vertices of  $G(\Pi)$ , and the *size* of  $\Pi$  is the sum of the sizes of the formulas that label its formula-vertices. Note that this definition of size does not depend on the weights that witness the proof. As we will see in the next section, such weights may be assumed to be integral and have small bit-complexity.

### 3.2 Checking the Global Condition

We still need to argue two facts about circular proofs: 1) that the existence of a witnessing flow assignment guarantees soundness, and 2) that its existence can be checked algorithmically in an efficient way. Soundness is proved in the next section. Here we argue that its existence can be checked efficiently. One way to do this is by solving a linear program.

**Lemma 1.** *There is a polynomial-time algorithm that, given as input a circular pre-proof  $\Pi$ , a finite set of hypothesis formulas  $\mathcal{H}$ , and a goal formula  $A$ , returns a flow assignment for  $\Pi$  that witnesses a proof of  $A$  from  $\mathcal{H}$ , if it exists.*

*Proof.* Let  $V = I \cup J$  be the set of vertices of the graph  $G(\Pi)$  of  $\Pi$ , partitioned into the set  $I$  of inference-vertices and the set  $J$  of formula-vertices. Observe that  $N^-(u) \subseteq I$  and  $N^+(u) \subseteq I$  for each  $u \in J$ . Let  $H \subseteq J$  be the set of formula-vertices whose labels are in  $\mathcal{H}$ , and let  $a \in J$  be a formula-vertex whose label is  $A$  and gets positive balance under some flow assignment for  $\Pi$ . For each  $w$  in  $I$ , let  $Y_w$  denote a real-valued variable and consider the following instance of the linear programming feasibility problem:

$$(P) : \begin{array}{ll} \sum_{w \in N^-(u)} Y_w - \sum_{w \in N^+(u)} Y_w \geq 1 & \text{for } u = a, \\ \sum_{w \in N^-(u)} Y_w - \sum_{w \in N^+(u)} Y_w \geq 0 & \text{for each } u \in J \setminus (H \cup \{a\}), \\ Y_w \geq 1 & \text{for each } w \in I. \end{array}$$

We claim that  $(P)$  has a feasible solution  $(y_w)_{w \in I}$  if and only if there exists a flow assignment  $F : I \rightarrow \mathbb{R}^+$  that witnesses a proof of  $A$  from  $\mathcal{H}$  by making the balance  $B(a)$  of  $a$  positive. For the *only if* direction, define  $F : I \rightarrow \mathbb{R}^+$  by  $F(w) := y_w$ , and read-off the required conditions for  $F$  from the inequalities that define  $(P)$ . For the *if* direction, define  $y_w := F(w)/D$ , where  $D$  is the minimum in the finite set  $\{F(w) : w \in I\} \cup \{B(a)\}$  and  $B(a)$  denotes the balance of  $a$ . Observe that  $D$  is strictly positive by definition and

the choice of  $a$ . The inequalities of  $(P)$  are satisfied by  $(y_w)_{w \in I}$  also by definition, and by the choice of  $D$ . Since the linear programming feasibility problem can be solved in polynomial time in the size of the input, the lemma follows.  $\square$

By elementary facts about linear programming (see [29]), it follows from this proof that if there is a flow assignment that witnesses a proof, then there is one with flows that are rational numbers whose bit-complexity is at most polynomial in the length of the circular pre-proof. By taking common denominators and multiplying through, the flows can even be taken to be positive integers of bit-complexity still polynomial in the length of the pre-proof. We collect these observations in a lemma.

**Lemma 2.** *Let  $\Pi$  be a circular pre-proof of length  $\ell$ . For every flow assignment  $F$  for  $\Pi$  there exists another flow assignment  $F'$  for  $\Pi$  such that:*

1.  $F'(w)$  is a positive integer bounded by  $\ell!$ , for every inference-vertex  $w$  of  $G(\Pi)$ ,
2.  $G(\Pi, F)$  and  $G(\Pi, F')$  have the same sets of sources and sinks.

*Proof.* Let  $I$  and  $J$  be the sets of inference- and formula-vertices of  $G(\Pi)$ . Let  $S \subseteq J$  and  $T \subseteq J$  be the sets of sources and sinks of  $G(\Pi, F)$ , respectively. Consider the following variant of the linear program  $(P)$  above:

$$(Q) : \begin{array}{ll} \sum_{w \in N^-(u)} Y_w - \sum_{w \in N^+(u)} Y_w \geq 1 & \text{for each } u \in T, \\ \sum_{w \in N^-(u)} Y_w - \sum_{w \in N^+(u)} Y_w \geq 0 & \text{for each } u \in J \setminus (S \cup T), \\ Y_w \geq 1 & \text{for each } w \in I. \end{array}$$

When we transform  $(Q)$  it into standard form by adding exactly  $|J| + |I| - |S|$  many slack variables, the result will be a linear program of the form  $Mx = b$ ,  $x \geq 0$  where  $x$  is a vector of  $2|I| + |J| - |S|$  variables,  $M$  is a constraint matrix of dimensions  $(|I| + |J| - |S|) \times (2|I| + |J| - |S|)$ , and  $b$  is a right-hand side  $(|I| + |J| - |S|)$ -vector. Moreover, each coefficient in the matrix  $M$  and the vector  $b$  will be in  $\{-1, 0, 1\}$ . Since this linear program has a solution (the one given by  $F$  adequately extended to the slack variables), it also has a basic feasible solution  $(x_u^*)_{u \in V}$ . Each component  $x_u^*$  is either 0 or, by Cramer's Rule, can be written in the form  $\det(N_u) / \det(N)$  where  $N$  is a square submatrix of  $M$ , and  $N_u$  is the matrix that results from replacing the column of  $N$  of index  $u$  by a subvector of the right-hand side vector  $b$ . By ignoring the slack variables we get a solution  $(y_w)_{w \in I}$  for  $(Q)$  of the same form. Multiplying through by the common denominator  $\det(N)$  we get an integral solution  $(y'_w)_{w \in I}$  for  $(Q)$  whose components have the form  $\det(N_w)$ ; none is 0 because  $Y_w \geq 1$  is one of the inequalities in  $(Q)$ . Each  $N_w$ -matrix has dimensions at most  $(|I| + |J| - |S|) \times (|I| + |J| - |S|)$ , and components in  $\{-1, 0, 1\}$ . It follows that  $y'_w = \det(N_w) \leq (|I| + |J| - |S|)! = \ell!$ . Taking  $F'(w) := y'_w$  for each  $w \in I$  completes the proof.  $\square$

### 3.3 Soundness of Circular Proofs

In this section we develop the soundness proof when the set of inference rules is fixed to axiom, symmetric cut, and split. See Section 2 for a discussion on this choice of rules. In the next section we discuss the general case.

We give two different proofs: one combinatorial and one (semi-)algebraic.

**Theorem 3.** *Let  $\mathcal{R}$  be the set of inference rules made of axiom, symmetric cut, and split. Let  $\mathcal{H}$  be a set of hypothesis formulas and let  $A$  be a goal formula. If there is a circular proof of  $A$  from  $\mathcal{H}$  through the rules in  $\mathcal{R}$ , then every truth assignment that satisfies every formula in  $\mathcal{H}$  also satisfies  $A$ .*

*First proof.* Fix a truth assignment  $\alpha$ . We prove the stronger claim that, for every circular pre-proof  $\Pi$  from an unspecified set of hypothesis formulas, every integral flow assignment  $F$  for  $\Pi$ , and every sink  $s$  of  $G(\Pi, F)$ , if  $\alpha$  falsifies the formula that labels  $s$ , then  $\alpha$  also falsifies the formula that labels some source of  $G(\Pi, F)$ . The proof is by induction on the the sum of the flows assigned by  $F$ , which we call the total flow-sum of  $F$ . Such induction is possible because we restrict to integral flow assignments, which is without loss of generality by Lemma 2.

If the total flow-sum is zero, then there are no inferences, hence there are no sinks, and the statement holds vacuously. Assume then that the total flow-sum is positive, and let  $s$  be a sink of  $G(\Pi, F)$ , with balance  $B(s) > 0$ , whose labelling formula  $B$  is falsified by  $\alpha$ . Since its balance is positive,  $s$  must have at least one in-neighbour  $r$ . Since the conclusion formula of the rule at  $r$  is falsified by  $\alpha$ , some premise formula of the rule at  $r$  must exist that is also falsified by  $\alpha$ . Let  $u$  be the corresponding in-neighbour of  $r$ , and let  $B(u)$  be its balance. If  $B(u)$  is negative, then  $u$  is a source of  $G(\Pi, F)$ , and we are done. Assume then that  $B(u)$  is non-negative.

Let  $\delta := \min\{B(s), F(r)\}$  and note that  $\delta > 0$  because  $B(s) > 0$  and  $F(r) > 0$ . We define a new circular pre-proof  $\Pi'$  and an integral flow assignment  $F'$  for  $\Pi'$  to which we will apply the induction hypothesis. The construction will guarantee the following properties:

1. the total flow-sum of  $F'$  is smaller than the total flow-sum of  $F$ .
2.  $u$  is a sink of  $G(\Pi', F')$  and  $s$  is not a source of  $G(\Pi', F')$ ,
3. if  $t$  is a source of  $G(\Pi', F')$ , then  $t$  is a source of  $G(\Pi, F)$  or an out-neighbour of  $r$  in  $G(\Pi)$ .

From this the claim will follow by applying the induction hypothesis to  $\Pi'$ ,  $F'$  and  $u$ . Indeed the induction hypothesis applies to them by Property 1 and the first half of Property 2. It will give a source  $t$  of  $G(\Pi', F')$  whose labelling formula is falsified by  $\alpha$ . We argue that  $t$  must also be a source of  $G(\Pi, F)$ , in which case we are done. To argue for this, assume otherwise and apply Property 3 to conclude that  $t$  is an out-neighbour of  $r$  in  $G(\Pi)$ , which by the second half of Property 2 must be different from  $s$  because  $t$  is a source of  $G(\Pi', F')$ . Recall now that  $s$  is a second out-neighbour of  $r$ . This can be the case only if  $r$  is a split inference, in which case the formulas that label  $s$  and  $t$  must be of the form  $C \vee B$  and  $C \vee \overline{B}$ , respectively, for appropriate formulas  $C$  and  $B$ . But, by assumption,  $\alpha$  falsifies the formula that labels  $s$ , namely  $C \vee B$ , which means that  $\alpha$  satisfies the formula  $C \vee \overline{B}$  that labels  $t$ . This is the contradiction we were after.

It remains to construct  $\Pi'$  and  $F'$  that satisfy Properties 1, 2, and 3. We define them by cases according to whether  $F(r) > B(s)$  or  $F(r) \leq B(s)$ , and then argue for the correctness of the construction. In case  $F(r) > B(s)$ , and hence  $\delta = B(s)$ , let  $\Pi'$  be defined as  $\Pi$  without change, and let  $F'$  be defined by  $F'(r) := F(r) - \delta$  and  $F'(w) := F(w)$  for every other  $w \in I \setminus \{r\}$ . Obviously  $\Pi'$  is still a pre-proof and  $F'$  is an integral flow assignment for  $\Pi'$  by the assumption that  $F(r) > B(s) = \delta$ . In case  $F(r) \leq B(s)$ , and hence  $\delta = F(r)$ , let  $\Pi'$  be defined as  $\Pi$  with the inference-step that labels  $r$  removed, and let  $F'$  be defined by  $F'(w) := F(w)$  for every  $w \in I \setminus \{r\}$ . Note that in this case  $\Pi'$  is still a pre-proof but perhaps from a larger set of hypothesis formulas.

In both cases the proof of the claim that  $\Pi'$  and  $F'$  satisfy Properties 1, 2, and 3 is the same. Property 1 follows from the fact that the total flow-sum of  $F'$  is the total flow-sum of  $F$  minus  $\delta$ , and  $\delta > 0$ . The first half of Property 2 follows from the fact that the balance of  $u$  in  $G(\Pi', F')$  is  $B(u) + \delta$ , while  $B(u) \geq 0$  by assumption and  $\delta > 0$ . The second half of Property 2 follows from the fact that the balance of  $s$  in  $G(\Pi', F')$  is  $B(s) - \delta$ , while  $B(s) \geq \delta$  by choice of  $\delta$ . Property 3 follows from the fact that the only formula-vertices

of  $G(\Pi', F')$  of balance smaller than that in  $G(\Pi, F)$  are the out-neighbours of  $r$ . This completes the proof of the claim, and of the theorem.  $\square$

We give a second, different proof of soundness that will play an important role later.

*Second proof.* Let  $\Pi$  be a circular pre-proof and let  $F$  be a flow assignment for  $\Pi$  that witnesses a proof of  $A$  from  $\mathcal{H}$ . Let  $\alpha$  be a truth assignment that satisfies all the formulas in  $\mathcal{H}$ , and let  $s$  be an arbitrary formula-vertex in  $G(\Pi)$ . We show that if  $\alpha$  falsifies the formula that labels  $s$ , then  $s$  is not a sink of  $G(\Pi, F)$ .

Let  $V = I \cup J$  be the set of vertices of  $G(\Pi)$  partitioned into the set  $I$  of inference-vertices, and the set  $J$  of formula-vertices. For every  $u \in J$ , let  $A_u$  be the formula that labels  $u$  and let  $Z_u := \alpha(A_u)$ ; the truth-value that  $\alpha$  gives to  $A_u$ . By inspection of the three allowed inference rules, for each  $w \in I$  with labelling inference rule  $R$  and in- and out-neighbours  $N^-$  and  $N^+$ , respectively, we have:

$$\begin{aligned} -(1 - Z_a) &\geq 0 && \text{if } R = \text{axiom with } N^+ = \{a\}, \\ (1 - Z_a) + (1 - Z_b) - (1 - Z_c) &\geq 0 && \text{if } R = \text{cut with } N^- = \{a, b\} \text{ and } N^+ = \{c\}, \\ (1 - Z_a) - (1 - Z_b) - (1 - Z_c) &\geq 0 && \text{if } R = \text{split with } N^- = \{a\} \text{ and } N^+ = \{b, c\}. \end{aligned}$$

Multiplying each such inequality by the positive flow  $F(w)$  of  $w$  and adding up over all  $w \in I$  we get

$$\sum_{w \in I} F(w) \left( \sum_{v \in N^-(w)} (1 - Z_v) - \sum_{u \in N^+(w)} (1 - Z_u) \right) \geq 0 \quad (7)$$

Rearranging the sum by formula-vertices, instead of arranging it by inference-vertices, we get

$$\sum_{u \in J} (1 - Z_u) \left( \sum_{w \in N^+(u)} F(w) - \sum_{w \in N^-(u)} F(w) \right) \geq 0. \quad (8)$$

The expression enclosed in parenthesis in the left-hand side in (8) equals  $-B(u)$ , where  $B(u)$  is the balance of  $u$  in  $G(\Pi, F)$ . Now,  $Z_u = 1$  whenever  $u$  is a source,  $Z_s = 0$  for  $s$  by assumption, and  $B(u)(1 - Z_u) \geq 0$  for every other formula-vertex  $u \in J$  by the definition of circular proof. Hence

$$-B(s) \geq 0, \quad (9)$$

which shows that  $s$  has non-positive balance in  $G(\Pi, F)$  and is thus not a sink.  $\square$

In the second proof of soundness, one can think of the  $Z_u$  as variables that are constrained by: 1) the inequalities that express the local soundness of the three types of inference rules, 2) the equations  $Z_u = 1$  for  $u$  a source, that express that each hypothesis is satisfied, and 3) the equation  $Z_s = 0$  for  $s$  a sink, that expresses that some conclusion is falsified. In other words, the constraints of types 1), 2) and 3) express that the proof is unsound, while equations (7) and (8) say that any valid flow assignment  $F(w)$  can play the role of a witness of its infeasibility.

### 3.4 Soundness for Other Sets of Rules

We claim that both proofs of soundness that we gave apply without change to any set of sound inference rules that have a single conclusion formula. This requirement is fulfilled by all sets of standard inference rules, such as (1), and is subsumed by the following more general but technical one:

(\*) Any inference rule in  $\mathcal{R}$  that has more than one conclusion formula has the property that any truth assignment that falsifies one of its conclusion formulas must satisfy all other conclusion formulas.

Obviously, if all rules in  $\mathcal{R}$  have a single conclusion, then (\*) is satisfied. Note also that the only rule that has more than one conclusion formula among axiom, symmetric cut, and split is split, and clearly it has the required property. Thus, the following statement generalizes Theorem 3.

**Theorem 4.** *Let  $\mathcal{R}$  be a set of sound inference rules that satisfy property (\*). Let  $\mathcal{H}$  be a set of hypothesis formulas and let  $A$  be a goal formula. If there is a circular proof of  $A$  from  $\mathcal{H}$  through the rules in  $\mathcal{R}$ , then every truth assignment that satisfies every formula in  $\mathcal{H}$  also satisfies  $A$ .*

*Proof.* The first proof of Theorem 3 was already phrased in a way that the generalization to sets of inference rules that satisfy (\*) is straightforward. We discuss the generalization of the second proof. Let  $\Pi$  be a circular proof with rules in  $\mathcal{R}$ , let  $A_u$  be the formula that labels the formula-vertex  $u$ , let  $Z_u := \alpha(A_u)$  be the truth value given to  $A_u$  by a truth assignment  $\alpha$ , and let  $w$  be an inference-vertex of  $\Pi$  with in- and out-neighbors  $N^-$  and  $N^+$ , respectively. Then, the following inequality holds:

$$\sum_{u \in N^-} (1 - Z_u) - \sum_{u \in N^+} (1 - Z_u) \geq 0 \quad (10)$$

Indeed, if  $Z_a = 0$  for some  $a \in N^+$ , then by the soundness of the rule there exists  $b \in N^-$  such that  $Z_b = 0$ , and by (\*) we also have  $Z_c = 1$  for every  $c \in N^+ \setminus \{a\}$ . The conclusion to this is that the left summand in (10) is at least 1, and the right summand in (10) is exactly 1, so their difference is non-negative. From here it suffices to note that this is the only property we used in order to derive equations (7), (8) and (9).  $\square$

## 4 Circular Resolution

In this section we investigate the power of Circular Resolution. Recall from the discussion in Section 2.4 that Resolution is traditionally defined to have cut as its only rule, but that an essentially equivalent version of it is obtained if we define it through symmetric cut, split, and axiom, still all restricted to clauses. This more liberal definition of Resolution, while staying equivalent vis-a-vis the tree-like and dag-like versions of Resolution, will play an important role for the circular version of Resolution.

While for Frege proof systems we will prove later that there is no qualitative difference between tree-like, dag-like, and circular proofs, in this section we show that circular Resolution can be exponentially stronger than dag-like Resolution. Indeed, we show that Circular Resolution is polynomially equivalent with the Sherali-Adams proof system, which is already known to be stronger than dag-like Resolution:

**Theorem 5.** *Sherali-Adams and Circular Resolution polynomially simulate each other. Moreover, the simulation one way converts monomial size  $s$  and degree  $d$  into size  $O(s)$  and width  $d$ , and the simulation in the reverse way converts size  $s$  and width  $w$  into monomial size  $O(s)$  and degree  $w$ .*

For the statement of Theorem 5 to even make sense, Sherali-Adams is to be understood as a proof system for deriving clauses from clauses, under an appropriate encoding of clauses as polynomial inequalities, to be discussed later in this section.

## 4.1 Pigeonhole Principles

We start by showing that the Pigeonhole Principle formula  $\text{PHP}_n^{n+1}$  has small Circular Resolution proofs. By the well-known lower bound of Haken [21], this will show that Circular Resolution is exponentially stronger than Resolution. However, we show the stronger claim that Circular Resolution is also stronger than Resolution when measured in terms of *width*; i.e., the length of the longest clause in the proof. To prove this, we need to introduce the *bipartite graph-based* variant of the Pigeonhole Principle from [6].

Let  $G$  be a bipartite graph with vertex bipartition  $(U, V)$ , and set of edges  $E \subseteq U \times V$ . For a vertex  $w \in U \cup V$ , we write  $N_G(w)$  to denote the set of neighbours of  $w$  in  $G$ , and  $\deg_G(w)$  to denote its degree. The Graph Pigeonhole Principle of  $G$ , denoted by  $G$ -PHP, is a CNF formula that has one variable  $X_{u,v}$  for each edge  $(u, v)$  in  $E$  and the following set clauses:

$$\begin{aligned} X_{u,v_1} \vee \cdots \vee X_{u,v_d} & \text{ for } u \in U \text{ with } N_G(u) = \{v_1, \dots, v_d\}, \\ \overline{X_{u_1,v}} \vee \overline{X_{u_2,v}} & \text{ for } u_1, u_2 \in U, v \in V \text{ with } u_1 \neq u_2, \text{ and } v \in N_G(u_1) \cap N_G(u_2). \end{aligned}$$

If  $|U| > |V|$ , and in particular if  $|U| = n + 1$  and  $|V| = n$ , then  $G$ -PHP is unsatisfiable by the pigeonhole principle. For  $G = K_{n+1,n}$ , the complete bipartite graph with sides of sizes  $n + 1$  and  $n$ , the formula  $G$ -PHP is the standard CNF encoding  $\text{PHP}_n^{n+1}$  of the pigeonhole principle.

Even for certain constant degree bipartite graphs with  $|U| = n + 1$  and  $|V| = n$ , the formulas are hard for Resolution.

**Theorem 6** ([6, 21]). *There are families of bipartite graphs  $(G_n)_{n \geq 1}$ , where  $G_n$  has maximum degree bounded by a constant and vertex bipartition  $(U, V)$  of  $G_n$  that satisfies  $|U| = n + 1$  and  $|V| = n$ , such that every Resolution refutation of  $G_n$ -PHP has width  $\Omega(n)$  and length  $2^{\Omega(n)}$ . Moreover, this implies that every Resolution refutation of  $\text{PHP}_n^{n+1}$  has length  $2^{\Omega(n)}$ .*

In contrast, we show that these formulas have Circular Resolution refutations of polynomial length and, simultaneously, constant width.

**Theorem 7.** *For every bipartite graph  $G$  of maximum degree  $d$  with bipartition  $(U, V)$  such that  $|U| > |V|$ , there is a Circular Resolution refutation of  $G$ -PHP of length polynomial in  $|U| + |V|$  and width  $d$ .*

*Proof.* We build the graph of the refutation in parts. Concretely, for every  $u \in U$  and  $v \in V$ , we describe two Circular Resolution proofs  $\Pi_{u \rightarrow}$  and  $\Pi_{\rightarrow v}$ , with their associated flow assignments. These proofs will have width bounded by  $\deg_G(u)$  and  $\deg_G(v)$ , respectively, and size polynomial in  $\deg_G(u)$  and  $\deg_G(v)$ , respectively. Moreover, the following properties will be ensured:

1. The proof-graph of  $\Pi_{u \rightarrow}$  contains a formula-vertex labelled by the empty clause 0 with balance +1 and a formula-vertex labelled  $\overline{X_{u,v}}$  with balance  $-1$  for every  $v \in N_G(u)$ ; any other formula-vertex that has negative balance is labelled by a clause of  $G$ -PHP.
2. The proof-graph  $\Pi_{\rightarrow v}$  contains a formula-vertex labelled by the empty clause 0 with balance  $-1$  and a formula-vertex labelled by  $\overline{X_{u,v}}$  with balance +1 for every  $u \in N_G(v)$ ; any other formula-vertex that has negative balance is labelled by a clause of  $G$ -PHP.




Figure 4: The diagram of the circular proof of  $\text{PHP}_3^4$ . The double circles indicate multiple inferences. The empty clause 0 is derived four times and used only three times.

By merging  $n$  of the  $n + 1$  many formula-vertices labelled by the empty clause in proofs of the form  $\Pi_{u \rightarrow}$  with the  $n$  many formula-vertices labelled by the empty clause in proofs of the form  $\Pi_{\rightarrow v}$  we get a proof in which all the formula-vertices that have negative balance are clauses of  $G$ -PHP, and the empty clause 0 has positive balance. This is indeed a Circular Resolution refutation of  $G$ -PHP. See Figure 4 for a diagram of the proof for  $\text{PHP}_3^4$ .

For the construction of  $\Pi_{u \rightarrow}$ , rename the neighbours of  $u$  as  $1, 2, \dots, \ell$ . Let  $C_j$  denote the clause  $X_{u,1} \vee \dots \vee X_{u,j}$  and note that  $C_\ell$  is a clause of  $G$ -PHP. Split  $\overline{X_{u,\ell}}$  on  $X_{u,1}$ , then split  $\overline{X_{u,\ell}} \vee X_{u,1}$  on  $X_{u,2}$ , then split  $\overline{X_{u,\ell}} \vee X_{u,1} \vee X_{u,2}$  on  $X_{u,3}$ , and so on until we produce  $\overline{X_{u,\ell}} \vee C_{\ell-1}$ . Then resolve this clause with  $C_\ell$  to produce  $C_{\ell-1}$ . As a sequence, this part of  $\Pi_{u \rightarrow}$  looks as follows:

$$\overline{X_{u,\ell}} \tag{11a}$$

$$\overline{X_{u,\ell}} \vee X_{u,1} \tag{11b}$$

$$\overline{X_{u,\ell}} \vee X_{u,1} \vee X_{u,2} \tag{11c}$$

$$\vdots \tag{11d}$$

$$\overline{X_{u,\ell}} \vee X_{u,1} \vee X_{u,2} \vee \dots \vee X_{u,\ell-1} \tag{11e}$$

$$X_{u,1} \vee X_{u,2} \vee \dots \vee X_{u,\ell-1}. \tag{11f}$$

We now repeat essentially the same construction: we start with  $\overline{X_{u,\ell-1}}$  and we split on  $X_{u,1}, X_{u,2}, \dots$ , as before until we produce  $\overline{X_{u,\ell-1}} \vee C_{\ell-2}$ . Cutting the latter with the previously deduced  $C_{\ell-1}$  gives  $C_{\ell-2}$ . We

continue other  $\ell - 2$  times in order to get down to the empty clause. Observe that we can set the flow of all splits and cuts in this proof to  $+1$  to get the balance claimed above.

For the construction of  $\Pi_{\rightarrow v}$  we need some more work. Again rename the neighbours of  $v$  as  $1, 2, \dots, \ell$ . All the inference steps in this construction will have assigned flow  $+1$ . The first step in building the proof  $\Pi_{\rightarrow v}$  is the derivation of the following sequence of clauses:

$$\overline{X_{1,v}} \tag{12a}$$

$$X_{1,v} \vee \overline{X_{2,v}} \tag{12b}$$

$$X_{1,v} \vee X_{2,v} \vee \overline{X_{3,v}} \tag{12c}$$

$$\vdots \tag{12d}$$

$$X_{1,v} \vee X_{2,v} \vee \dots \vee \overline{X_{\ell-1,v}} \tag{12e}$$

$$X_{1,v} \vee X_{2,v} \vee \dots \vee X_{\ell-1,v}. \tag{12f}$$

Notice that (12f) does not follow the pattern of the previous clauses. We start these inferences by splitting the empty clause on variable  $X_{1,v}$ , to get clauses  $\overline{X_{1,v}}$  and  $X_{1,v}$ . We split the latter on  $X_{2,v}$  to get  $X_{1,v} \vee \overline{X_{2,v}}$  and  $X_{1,v} \vee X_{2,v}$ . For  $i = 3, \dots, \ell - 1$  we keep splitting  $X_{1,v} \vee \dots \vee X_{i-1,v}$  on  $X_{i,v}$  to get the clauses  $X_{1,v} \vee \dots \vee X_{i-1,v} \vee \overline{X_{i,v}}$  and  $X_{1,v} \vee \dots \vee X_{i-1,v} \vee X_{i,v}$ . The empty clause has balance  $-1$  and all clauses (12a–12f) have balance  $+1$ .

Now to complete the proof  $\Pi_{\rightarrow v}$  we deduce the  $\ell$  singleton clauses  $\overline{X_{1,v}}, \dots, \overline{X_{\ell,v}}$  from the clauses (12a–12f). Observe that a simple sequence of cut rules between a clause  $X_{1,v} \vee \dots \vee X_{i-1,v} \vee \overline{X_{i,v}}$  and clauses  $\overline{X_{j,v}} \vee \overline{X_{i,v}}$  for  $1 \leq j < i$  produces the singleton clause  $\overline{X_{i,v}}$ , furthermore its balance is  $+1$ . We do a similar sequence of steps using  $X_{1,v} \vee \dots \vee X_{\ell-2,v} \vee X_{\ell-1,v}$  and clauses  $\overline{X_{j,v}} \vee \overline{X_{\ell,v}}$  for  $1 \leq j < \ell$  to produce the clause  $\overline{X_{\ell,v}}$  with balance  $+1$ , and conclude the construction of  $\Pi_{\rightarrow v}$ .  $\square$

We get the following immediate consequence:

**Corollary 8.** *The pigeonhole formulas  $\text{PHP}_n^{n+1}$  have Circular Resolution refutations of polynomial length.*

In combination with Theorem 6, this shows that Circular Resolution can be exponentially stronger than Resolution. In the next two sections we determine the exact power of Circular Resolution.

## 4.2 Simulation by Sherali-Adams

In this section we prove one half of Theorem 5. We need some preparation. Fix a set of variables  $X_1, \dots, X_n$  and their twins  $\overline{X}_1, \dots, \overline{X}_n$ . For a clause  $C = \bigvee_{j \in Y} X_j \vee \bigvee_{j \in Z} \overline{X}_j$ , define

$$T(C) := - \prod_{j \in Y} \overline{X}_j \prod_{j \in Z} X_j, \tag{13}$$

Observe that a truth assignment satisfies  $C$  if and only if the corresponding 0-1 assignment for the variables of  $T(C)$  makes the inequality  $T(C) \geq 0$  true. There is an alternative encoding of clauses into inequalities that is sometimes used. Define

$$L(C) := \sum_{j \in Y} X_j + \sum_{j \in Z} \overline{X}_j - 1, \tag{14}$$

and observe that a truth assignment satisfies  $C$  if and only if the corresponding 0-1 assignment makes the inequality  $L(C) \geq 0$  true. We state the results of this section for the  $T$ -encoding of clauses, but the same result would hold for the  $L$ -encoding because there is an efficient SA proof of (13) from (14) (see Lemma 4.2 in [3]), and vice-versa.

We will use the following lemma, which is a variant of Lemma 4.4 in [3]:

**Lemma 9.** *Let  $w \geq 2$  be an integer, let  $C$  be a clause with at most  $w$  literals, let  $D$  be a clause with at most  $w - 1$  literals, and let  $X$  be a variable that does not appear in  $D$ . Then the following four inequalities have Sherali-Adams proofs (from nothing) of constant monomial size and degree  $w$ :*

1.  $T(X \vee \bar{X}) \geq 0$ ,
2.  $-T(D \vee \bar{X}) - T(D \vee X) + T(D) \geq 0$ ,
3.  $-T(D) + T(D \vee \bar{X}) + T(D \vee X) \geq 0$ ,
4.  $-T(C) \geq 0$ .

*Proof.* Let  $D = \bigvee_{i \in Y} X_i \vee \bigvee_{j \in Z} X_j$  and  $C = \bigvee_{i \in Y'} X_i \vee \bigvee_{j \in Z'} X_j$ . Then

1.  $T(X \vee \bar{X}) = (1 - X - \bar{X}) \cdot X + (X^2 - X)$ ,
2.  $-T(D \vee \bar{X}) - T(D \vee X) + T(D) = (X + \bar{X} - 1) \cdot \prod_{i \in Y} \bar{X}_i \prod_{j \in Z} X_j$ ,
3.  $-T(D) + T(D \vee \bar{X}) + T(D \vee X) = (1 - X - \bar{X}) \cdot \prod_{i \in Y} \bar{X}_i \prod_{j \in Z} X_j$ ,
4.  $-T(C) = 1 \cdot \prod_{i \in Y'} \bar{X}_i \prod_{j \in Z'} X_j$ .

The claim on the monomial size and the degree follows. □

Now we are ready to state and prove the first half of Theorem 5.

**Lemma 10.** *Let  $A_1, \dots, A_m$  and  $A$  be clauses. If there is a Circular Resolution proof of  $A$  from  $A_1, \dots, A_m$  of length  $s$  and width  $w$ , then there is a Sherali-Adams proof of  $T(A) \geq 0$  from  $T(A_1) \geq 0, \dots, T(A_m) \geq 0$  of monomial size  $O(s)$  and degree  $w$ .*

*Proof.* Let  $\Pi$  be a Circular Resolution proof of  $A$  from  $A_1, \dots, A_m$ , and let  $F$  be the corresponding flow assignment. Let  $I$  and  $J$  be the sets of inference- and formula-vertices of  $G(\Pi)$ , and let  $B(u)$  denote the balance of formula-vertex  $u \in J$  in  $G(\Pi, F)$ . For each formula-vertex  $u \in J$  labelled by formula  $A_u$ , define the polynomial  $P_u := T(A_u)$ . For each inference-vertex  $w \in I$  labelled by rule  $R$ , with sets of in- and out-neighbours  $N^-$  and  $N^+$ , respectively, define the polynomial

$$\begin{aligned} P_w &:= T(A_a) && \text{if } R = \text{axiom with } N^+ = \{a\}, \\ P_w &:= -T(A_a) - T(A_b) + T(A_c) && \text{if } R = \text{cut with } N^- = \{a, b\} \text{ and } N^+ = \{c\}, \\ P_w &:= -T(A_a) + T(A_b) + T(A_c) && \text{if } R = \text{split with } N^- = \{a\} \text{ and } N^+ = \{b, c\}. \end{aligned}$$

By double counting, the following polynomial identity holds:

$$\sum_{u \in J} B(u) P_u = \sum_{w \in I} F(w) P_w. \tag{15}$$

By hypothesis  $G(\Pi, F)$  has a sink  $s$  labelled by the derived clause  $A$ . Since  $B(s) > 0$ , equation (15) rewrites into

$$\sum_{w \in I} \frac{F(w)}{B(s)} P_w + \sum_{u \in J \setminus \{s\}} -\frac{B(u)}{B(s)} P_u = P_s.$$

We claim that this identity is a legitimate Sherali-Adams proof of  $T(A) \geq 0$  from the inequalities  $T(A_1) \geq 0, \dots, T(A_m) \geq 0$ . First,  $P_s = T(A_s) = T(A)$ , i.e., the right-hand side is correct. Second, each term  $(F(w)/B(s))P_w$  for  $w \in I$  is a sum of legitimate terms of a Sherali-Adams proof by the definition of  $P_w$  and Parts 1, 2 and 3 of Lemma 9. Third, since each source  $u \in I$  of  $G(\Pi, F)$  has  $B(u) < 0$  and is labelled by a formula in  $A_1, \dots, A_m$ , the term  $(-B(u)/B(s))P_u$  of a source  $u \in I$  is a positive multiple of  $T(A_u)$  and hence also a legitimate term of a Sherali-Adams proof from  $T(A_1) \geq 0, \dots, T(A_m) \geq 0$ . And fourth, since each non-source  $u \in I$  of  $G(\Pi, F)$  has  $B(u) \geq 0$ , each term  $(-B(u)/B(s))P_u$  of a non-source  $u \in I$  is a sum of legitimate terms of a Sherali-Adams proof by the definition of  $P_u$  and Part 4 of Lemma 9. The monomial size and degree of this Sherali-Adams proof are as claimed, and the proof of the Lemma is complete.  $\square$

### 4.3 Simulation of Sherali-Adams

In this section we prove the other half of Theorem 5. We use the notation from Section 4.2.

**Lemma 11.** *Let  $A_1, \dots, A_m$  and  $A$  be non-tautological clauses. If there is a Sherali-Adams proof of  $T(A) \geq 0$  from  $T(A_1) \geq 0, \dots, T(A_m) \geq 0$  of monomial size  $s$  and degree  $d$ , then there is a Circular Resolution proof of  $A$  from  $A_1, \dots, A_m$  of length  $O(s)$  and width  $d$ .*

*Proof.* Fix a Sherali-Adams proof of  $T(A) \geq 0$  from  $T(A_1) \geq 0, \dots, T(A_m) \geq 0$ , say

$$\sum_{j=1}^t Q_j P_j = T(A), \tag{16}$$

where  $Q_j$  is a non-negative linear combination of monomials on the variables  $X_1, \dots, X_n$  and  $\bar{X}_1, \dots, \bar{X}_n$ , and  $P_j$  is a polynomial from among  $T(A_1), \dots, T(A_m)$  or from among the polynomials in the list (5) from the definition of Sherali-Adams in Section 2.

Our goal is to massage the proof (16) until it becomes a Circular Resolution proof in disguise. Towards this, as a first step, we claim that (16) can be transformed into a *normalized proof* of the form

$$\sum_{j=1}^{t'} Q'_j P'_j = T(A) \tag{17}$$

that has the following properties:

1. each  $Q'_j$  is a positive multiple of a multilinear monomial, and  $Q'_j P'_j$  is multilinear,
2. each  $P'_j$  is a polynomial among  $T(A_1), \dots, T(A_m)$ , or among the polynomials in the set

$$\{-X_i \bar{X}_i, 1 - X_i - \bar{X}_i, X_i + \bar{X}_i - 1 : i \in [n]\} \cup \{1\}. \tag{18}$$

Comparing (18) with the original list (5) in the definition of Sherali-Adams, note that we have replaced the polynomials  $X_i - X_i^2$  and  $X_i^2 - X_i$  by  $-X_i\bar{X}_i$ . Note also that, by splitting the  $Q_j$ 's into their terms, we may assume without loss of generality that each  $Q_j$  in (16) is a positive multiple of a monomial on the variables  $X_1, \dots, X_n$  and  $\bar{X}_1, \dots, \bar{X}_n$ .

In order to prove the claim we rely on the well-known fact that each real-valued function over the Boolean domain has a unique representation as a multilinear polynomial:

**Fact 12.** *For every natural number  $N$  and every function  $f : \{0, 1\}^N \rightarrow \mathbb{R}$  there is a unique multilinear polynomial  $P$  with  $N$  variables satisfying  $P(a_1, \dots, a_N) = f(a_1, \dots, a_N)$  for every  $a_1, \dots, a_N \in \{0, 1\}$ .*

With this fact in hand, it suffices to convert each  $Q_j P_j$  in the left-hand side of (16) into a  $Q'_j P'_j$  of the required form (or 0), and check that  $Q_j P_j$  and  $Q'_j P'_j$  are equivalent over the 0-1 assignments to its variables (without relying on the constraint that  $\bar{X}_i = 1 - X_i$ ). The claim will follow from the combination of Fact 12 and the fact that  $T(A)$  is multilinear since, by assumption,  $A$  is non-tautological.

We proceed to the conversion of each  $Q_j P_j$  into a  $Q'_j P'_j$  of the required form. Recall that we assumed already, without loss of generality, that each  $Q_j$  is a positive multiple of a monomial. The multilinearization of a monomial  $Q_j$  is the monomial  $M(Q_j)$  that results from replacing every factor  $Y^k$  with  $k \geq 2$  in  $Q_j$  by  $Y$ . Obviously  $Q_j$  and  $M(Q_j)$  agree on 0-1 assignments, but replacing each  $Q_j$  by  $M(Q_j)$  is not enough to guarantee the normal form that we are after. We need to proceed by cases on  $P_j$ .

If  $P_j$  is one of the polynomials among  $T(A_1), \dots, T(A_m)$ , say  $T(A_i)$ , then we let  $Q'_j$  be  $M(Q_j)$  with every variable that appears in  $A_i$  deleted, and let  $P'_j$  be  $T(A_i)$  itself. It is obvious that this works. If  $P_j$  is  $1 - X_i - \bar{X}_i$ , then we proceed by cases on whether  $Q_j$  contains  $X_i$  or  $\bar{X}_i$  or both or neither. If  $Q_j$  contains neither  $X_i$  nor  $\bar{X}_i$ , then the choice  $Q'_j = M(Q_j)$  and  $P'_j = P_j$  works. If  $Q_j$  contains  $X_i$  or  $\bar{X}_i$ , call it  $Y$ , but not both, then the choice  $Q'_j = M(Q_j)/Y$  and  $P'_j = -X_i\bar{X}_i$  works. If  $Q_j$  contains both  $X_i$  and  $\bar{X}_i$ , then the choice  $Q'_j = M(Q_j)/(X_i\bar{X}_i)$  and  $P'_j = -X_i\bar{X}_i$  works. If  $P_j$  is  $X_i + \bar{X}_i - 1$ , then again we proceed by cases on whether  $Q_j$  contains  $X_i$  or  $\bar{X}_i$  or both or neither. If  $Q_j$  contains neither  $X_i$  nor  $\bar{X}_i$ , then the choice  $Q'_j = M(Q_j)$  and  $P'_j = P_j$  works. If  $Q_j$  contains  $X_i$  or  $\bar{X}_i$ , call it  $Y$ , but not both, then the choice  $Q'_j = M(Q_j)\bar{Y}$  and  $P'_j = 1$  works. If  $Q_j$  contains both  $X_i$  and  $\bar{X}_i$ , then the choice  $Q'_j = M(Q_j)$  and  $P'_j = 1$  works. If  $P_j$  is the polynomial 1, then the choice  $Q'_j = M(Q_j)$  and  $P'_j = 1$  works. Finally, if  $P_j$  is of the form  $X_i - X_i^2$  or  $X_i^2 - X_i$ , then we replace  $Q_j P_j$  by 0. Observe that in this case  $Q_j P_j$  is always 0 over 0-1 assignments, and the conversion is correct. This completes the proof that (17) exists.

It remains to be seen that the normalized proof (17) is a Circular Resolution proof in disguise. For each  $j \in [t']$ , let  $a_j$  and  $M_j$  be the positive real and the multilinear monomial, respectively, such that  $Q'_j = a_j M_j$ . Let also  $C_j$  be the unique clause on the variables  $X_1, \dots, X_n$  such that  $T(C_j) = -M_j$ . Let  $[t']$  be partitioned into five sets  $I_0 \cup I_1 \cup I_2 \cup I_3 \cup I_4$  where

1.  $I_0$  is the set of  $j \in [t']$  such that  $P'_j = T(A_{i_j})$  for some  $i_j \in [m]$ ,
2.  $I_1$  is the set of  $j \in [t']$  such that  $P'_j = -X_{i_j}\bar{X}_{i_j}$  for some  $i_j \in [n]$ ,
3.  $I_2$  is the set of  $j \in [t']$  such that  $P'_j = 1 - X_{i_j} - \bar{X}_{i_j}$  for some  $i_j \in [n]$ ,
4.  $I_3$  is the set of  $j \in [t']$  such that  $P'_j = X_{i_j} + \bar{X}_{i_j} - 1$  for some  $i_j \in [n]$ ,
5.  $I_4$  is the set of  $j \in [t']$  such that  $P'_j = 1$ .

Note that the pair of sets  $I_0$  and  $I_1$  is disjoint because each clause  $A_{i_j}$  is non-tautological by assumption. Since the rest of pairs are clearly disjoint, we get a partition of  $[t']$ . Define new polynomials  $P_j''$  as follows:

$$\begin{aligned} P_j'' &:= T(C_j \vee A_{i_j}) \text{ for } j \in I_0, \\ P_j'' &:= T(C_j \vee \overline{X_{i_j}} \vee X_{i_j}) \text{ for } j \in I_1, \\ P_j'' &:= -T(C_j) + T(C_j \vee \overline{X_{i_j}}) + T(C_j \vee X_{i_j}) \text{ for } j \in I_2, \\ P_j'' &:= -T(C_j \vee \overline{X_{i_j}}) - T(C_j \vee X_{i_j}) + T(C_j) \text{ for } j \in I_3, \\ P_j'' &:= T(C_j) \text{ for } j \in I_4. \end{aligned}$$

With this notation, (17) rewrites into

$$\sum_{j \in I_0} a_j P_j'' + \sum_{j \in I_1} a_j P_j'' + \sum_{j \in I_2} a_j P_j'' + \sum_{j \in I_3} a_j P_j'' = T(A) + \sum_{j \in I_4} a_j P_j''. \quad (19)$$

Finally we are ready to construct the circular proof. We build it by listing the inference-vertices with their associated flows, and then we identify together all the formula-vertices that are labelled by the same clause.

Intuitively,  $I_0$ 's are weakenings of hypothesis clauses,  $I_1$ 's are weakenings of axioms,  $I_2$ 's are cuts,  $I_3$ 's are splits, and  $I_4$ 's can be thought of as left-overs, i.e., clauses that were produced but never used. Formally, each  $j \in I_0$  becomes a chain of  $|C_j|$  many split vertices that starts at the hypothesis clause  $A_{i_j}$  and produces its weakening  $C_j \vee A_{i_j}$ ; all split vertices in this chain have flow  $a_j$ . Each  $j \in I_1$  becomes a sequence that starts at one axiom vertex that produces  $X_{i_j} \vee \overline{X_{i_j}}$  with flow  $a_j$ , followed by a chain of  $|C_j|$  many split vertices that produces its weakening  $C_j \vee X_{i_j} \vee \overline{X_{i_j}}$ ; all split vertices in this chain also have flow  $a_j$ . Each  $j \in I_2$  becomes one cut vertex that produces  $C_j$  from  $C_j \vee X_{i_j}$  and  $C_j \vee \overline{X_{i_j}}$  with flow  $a_j$ . And each  $j \in I_3$  becomes one split vertex that produces  $C_j \vee X_{i_j}$  and  $C_j \vee \overline{X_{i_j}}$  from  $C_j$  with flow  $a_j$ . Note that some of the produced clauses, such as the second conclusion in certain intermediate splits, may never be used as hypothesis in other rules.

This defines the inference-vertices of the proof graph. The construction is completed by introducing one formula-vertex for each different clause that occurs as a premise or a conclusion of these inference-vertices. The construction was designed in such a way that equation (19) witnesses that this proof graph and its associated flow assignment makes a correct circular proof. In order to see this we need to compute the balances of all the formula-vertices, check that some sink is labelled  $A$ , and check that all sources are labelled by formulas in  $A_1, \dots, A_m$ .

First note that, for each formula-vertex  $u$  whose labelling formula appears as an intermediate formula in a chain of split inferences for some  $j \in I_0 \cup I_1$ , the contribution of this occurrence to the balance of  $u$  is zero. Indeed, for being an intermediate formula in the chain, the flow  $a_j$  of the inference that produces it cancels out the flow  $a_j$  of the inference that consumes it. Second note that the only formula-vertices  $u$  whose labelling formulas are of the form  $\overline{X_{i_j}} \vee X_{i_j}$  are those that are introduced by an axiom inference in case  $j \in I_1$ ; this follows from the multilinearity of the  $Q_j' P_j'$  and the assumption that the  $A_i$  are non-tautological. It follows that the balance of such  $u$  is positive; indeed its balance is the flow  $a_j$  of the axiom inference that produces it. Third note that we do not need to compute the balance of the formula-vertices  $u$  whose labelling formula is a hypothesis among  $A_1, \dots, A_m$ ; any balance is allowed for it. With these three notes in mind it suffices to check that:

1. some sink is labelled  $A$ ,
2. for  $j \in I_0$  the formula-vertex for  $C_j \vee A_{i_j}$  has non-negative balance,
3. for  $j \in I_1$  the formula-vertex for  $C_j \vee \overline{X_{i_j}} \vee X_{i_j}$  has non-negative balance,
4. for  $j \in I_2 \cup I_3$ , the vertices for  $C_j$ ,  $C_j \vee \overline{X_{i_j}}$ , and  $C_j \vee X_{i_j}$  have non-negative balance.

For (1) it suffices to note that  $T(A)$  appears in the right-hand side of (19) with a positive multiplier, hence its occurrences in the left-hand side must have multipliers that add up to a positive quantity. For (2), (3), and (4) it suffices to note that if  $B$  is one of the indicated formulas, then  $T(B)$  appears in the left-hand side of (19) with multipliers that add-up to either a positive quantity or 0 depending on whether  $T(B)$  appears in the right-hand side or not, respectively. Hence its balance is non-negative, and the proof that the circular proof is correct is complete. The claim that the length of this proof is  $O(s)$  and its width is  $d$  follows by inspection.  $\square$

## 5 Circular Frege vs Tree-Like Frege

For some weak proof systems, such as Resolution, it makes a great deal of difference whether the proof-graph has tree-like structure or not [8]. For stronger proofs systems, such as Frege, this is not the case. Indeed Tree-like Frege polynomially simulates Dag-like Frege, and this holds true of any inference-based proof system with the set of all formulas as its set of allowed formulas, and a finite set of inference rules that is implicationally complete [23]. Since circular proofs further generalize the structure of the proof-graph, it is interesting to discuss whether circular proofs in Frege are complexity-wise more powerful than standard Frege proofs.

It turns out that this is not the case. In this section we show how to efficiently simulate Circular Frege, as defined in Section 3, by standard Frege proofs.

**Theorem 13.** *Tree-like Frege and Circular Frege polynomially simulate each other.*

The main idea underlying the simulation of Circular Frege by standard Frege is to formalize, in standard Dag-like Frege itself, the LP-based proof of soundness of Frege circular proofs; cf. the *second proof* of Theorem 3. To do that we use a formalization of linear arithmetic in Frege, due to Buss [12] and Goerdt [19]. This formalization was originally designed to simulate counting arguments and Cutting Planes in Frege. Since Cutting Planes subsumes LP-reasoning, the core of the LP-based proof of Theorem 3 can be formalized in it. It should be pointed out that, while the Frege systems that were used by Buss [12] and Goerdt [19] are not exactly the same as ours, their results apply to our definition of Frege; this follows from the well-known robustness properties of Frege systems which guarantee that any two (Dag-like) Frege systems polynomially simulate each other [13].

### 5.1 Formalization of Linear Arithmetic in Frege

We collect the relevant parts of Goerdt's results in a single theorem. Before that, we need to introduce some notation. Fix a set of  $n$  Boolean variables  $X_1, \dots, X_n$ . Let  $\mathcal{L}$  denote the collection of all linear inequalities of the form

$$a_1 X_1 + \dots + a_n X_n \geq b, \tag{20}$$

where  $X_1, \dots, X_n$  are formal variables and  $a_1, \dots, a_n$  and  $b$  are integers. Our notation includes inequalities of the form  $0 \geq b$ , and  $a_{i_1}X_{i_1} + \dots + a_{i_t}X_{i_t} \geq b$  by deleting the terms with zero coefficients. By the *set of variables* of an inequality we mean the set of variables that appear in it with non-zero coefficient.

Let  $\ell$  and  $\ell'$  denote two inequalities in  $\mathcal{L}$ , with sequences of coefficients  $b, a_1, \dots, a_n$  and  $b', a'_1, \dots, a'_n$ . Let  $c$  be a positive integer. We write  $c \cdot \ell$  and  $\ell + \ell'$  for the following two inequalities, respectively:

$$\begin{aligned} ca_1X_1 + \dots + ca_nX_n &\geq cb, \\ (a_1 + a'_1)X_1 + \dots + (a_n + a'_n)X_n &\geq (b + b'). \end{aligned}$$

Let  $\mathcal{F}$  denote the collection of all propositional formulas in negation normal form. We move back and forth between the truth assignments and the 0-1 assignments for the same sets of variables. If  $f : \{X_1, \dots, X_n\} \rightarrow \{0, 1\}$  denotes such an assignment, and  $\ell$  and  $A$  denote, respectively, an inequality and a formula on the variables  $X_1, \dots, X_n$ , then we write  $f(\ell)$  and  $f(A)$  for their truth values under  $f$ . Concretely, if  $\ell$  is as in (20), then  $f(\ell)$  is true if and only if  $a_1f(X_1) + \dots + a_nf(X_n)$  is at least  $b$ .

For concreteness, we specify a size measure for inequalities. An inequality  $\ell$  as in (20) is represented by the sequence of the binary encodings of its coefficients  $b, a_1, \dots, a_n$ ; its size is  $\Theta(n + \log_2(|b|) + \sum_{i=1}^n \log_2(|a_i|))$ , with the convention that  $\log_2(0) = 0$ , and where  $|v|$  denotes the absolute value of  $v \in \mathbb{Z}$ .

**Theorem 14** ([19]). *There is a mapping  $I : \mathcal{L} \rightarrow \mathcal{F}$  that takes linear inequalities to formulas and that has the following properties. For every two inequalities  $\ell$  and  $\ell'$  in  $\mathcal{L}$ , every positive integer  $c$ , every truth assignment  $f$ , and every variable  $X$ , the following hold:*

1.  $I(\ell)$  has size polynomial in the size of  $\ell$ ,
2.  $I(\ell)$  has the same variables as  $\ell$ , and  $f(\ell) = f(I(\ell))$ ,
3. there is a polynomial-size Frege proof of  $I(\ell + \ell')$  from  $I(\ell)$  and  $I(\ell')$ ,
4. there is a polynomial-size Frege proof of  $I(c \cdot \ell)$  from  $I(\ell)$ ,
5. there is a polynomial-size Frege proof of  $I(\ell)$  from  $I(c \cdot \ell)$ ,
6. there is a polynomial-size Frege proof of  $I(X \geq 1)$  from  $X$ ,
7. there is a polynomial-size Frege proof of  $I(-X \geq 0)$  from  $\overline{X}$ ,
8. there is a polynomial-size Frege proof of  $I(-X \geq -1)$  from nothing,
9. there is a polynomial-size Frege proof of  $I(X \geq 0)$  from nothing,
10. there is a polynomial-size Frege proof of  $I(0 \geq 0)$  from nothing,
11. there is a polynomial-size Frege proof of  $0$  from  $I(0 \geq 1)$ .

Moreover, the mapping  $I$  and the Frege proofs in Points 3–11 are all computable in time that is bounded by a fixed polynomial in the sizes of the input and the output inequalities.

*Proof.* All this can be found in Goerd't's article [19], which in turn builds on Buss's seminal [12]: the definition of the mapping  $I$  is in Section 2.6 of Goerd't's article, Points 1–2, and Points 6–11 follow by inspection of the definition of  $I$  given there, and Points 3, 4, and 5 are Theorems 3.1, 3.5, and 3.6 in Goerd't's article, respectively.  $\square$

Technically the inequalities are defined in (20) with the variables on the left side of the  $\geq$  relation symbol and the constant on the right side. For readability reasons, in the following we apply  $I$  to linear inequalities written in any form that follows from adding or subtracting the same linear terms from both sides of an inequality in official form; e.g., when we write  $I(-1 \geq 0)$  or  $I(X_1 - 2 \geq X_2)$  we really mean  $I(0 \geq 1)$  and  $I(X_1 - X_2 \geq 2)$ , respectively.

## 5.2 Proof of the Simulation

This section is devoted to the proof of Theorem 4. The statement that Circular Frege polynomially simulates Tree-like Frege follows from the discussion in Section 2.4. We concentrate on the reverse simulation. Since it is known that Tree-like Frege polynomially simulates Dag-like Frege, it suffices to do the simulation through dag-like proofs. Also we claim that it suffices to do the simulation only for refutations. Indeed, from a short circular proof of  $A$  from  $\mathcal{H}$  we can get a short circular refutation of  $\mathcal{H} \cup \{\overline{A}\}$  by adding a cut between the derived  $A$  and the new hypothesis  $\overline{A}$ , with flow equal to the balance of the formula-vertex of  $A$ . And from a short dag-like refutation of  $\mathcal{H} \cup \{\overline{A}\}$  we can get a short dag-like proof of  $A$  from  $\mathcal{H}$  by replacing each use of the hypothesis formula  $\overline{A}$  by the axiom instance  $A \vee \overline{A}$ .

Let  $\Pi$  be a Circular Frege refutation of a set of hypothesis formulas  $\mathcal{H}$ . Let us choose an arbitrary  $s$  among the formula-vertices in  $G(\Pi)$  that are labelled by the empty formula. The simulation goes in three steps. In the first step we build an infeasible linear program  $P = \{\ell_1, \dots, \ell_m\}$  that has one variable  $Z_u$  for each formula-vertex  $u$  of  $G(\Pi)$ . Any witness of infeasibility of  $P$  will witness the soundness of  $\Pi$  as a circular refutation. This is done by closely following the second proof of soundness of circular proofs; cf. Theorem 3. In the second step we apply Theorem 14 to convert an LP-based witness of infeasibility for  $P$  into a Frege refutation of the set of formulas  $\mathcal{H}' := \{I(\ell_1), \dots, I(\ell_m)\}$ . Here  $I$  is the mapping from Theorem 14. In the third step we apply the substitution defined by  $Z_u := A_u$  to this Frege refutation, where  $A_u$  is the formula that labels the formula-vertex  $u$ , and we apply Theorem 14 again to show that each formula in the substituted  $\mathcal{H}'$  has an efficient Frege proof from  $\mathcal{H}$ .

*First step.* The linear program  $P$  has one variable  $Z_u$  for each formula-vertex  $u \in J$  in  $G(\Pi)$ , and two sets of inequalities  $P_J = \{\ell_u : u \in J\}$  and  $P_I = \{\ell_w : w \in I\}$  indexed by the sets of formula-vertices  $J$  and inference-vertices  $I$  of  $G(\Pi)$ , respectively. Concretely, for each formula-vertex  $u \in J$  the inequality  $\ell_u$  is defined as follows:

$$\begin{aligned} -Z_s &\geq 0 && \text{for the formula-vertex } s \text{ labelled by the derived empty formula,} \\ -(1 - Z_u) &\geq 0 && \text{if } u \text{ is a formula-vertex of a hypothesis formula,} \\ (1 - Z_u) &\geq 0 && \text{if } u \text{ is any other formula-vertex.} \end{aligned}$$

For each inference-vertex  $w$  with labelling inference rule  $R$  and in- and out-neighbours  $N^-$  and  $N^+$ , respectively, the inequality  $\ell_w$  is defined as follows:

$$\begin{aligned} -(1 - Z_a) &\geq 0 && \text{if } R = \text{axiom with } N^+ = \{a\}, \\ (1 - Z_a) + (1 - Z_b) - (1 - Z_c) &\geq 0 && \text{if } R = \text{cut with } N^- = \{a, b\} \text{ and } N^+ = \{c\}, \\ (1 - Z_a) - (1 - Z_b) - (1 - Z_c) &\geq 0 && \text{if } R = \text{split with } N^- = \{a\} \text{ and } N^+ = \{b, c\}. \end{aligned}$$

A certificate of the infeasibility of  $P = P_I \cup P_J$  is given by two assignments of non-negative weights ( $b_u : u \in J$ ) and ( $c_w : w \in I$ ) for the inequalities in  $P_J$  and  $P_I$ , respectively, in such a way that the corresponding positive linear combination

$$\sum_{u \in J} b_u \cdot \ell_u + \sum_{w \in I} c_w \cdot \ell_w \tag{21}$$

simplifies to the trivially false inequality  $-1 \geq 0$  (or equivalently  $0 \geq 1$ ). In turn, such an assignment of weights can be shown to exist from the assumption that  $\Pi$  is a valid circular refutation: let  $F$  be a flow assignment that witnesses that  $\Pi$  is a valid proof and  $B$  the corresponding balance on the formula-vertices, and set  $b_u := -B(u)/B(s)$  for each formula-vertex  $u \in J$  that is a source of  $G(\Pi, F)$ , set  $b_u := B(u)/B(s)$  for each formula-vertex  $u \in J$  that is not a source of  $G(\Pi, F)$ , and set  $c_w := F(w)/B(s)$  for every inference-vertex  $w \in I$ . Note that  $B(s)$  is strictly positive because  $s$  must be a sink of  $G(\Pi, F)$ . This means that each  $b_u$  is well-defined and non-negative because the balance of all formula-vertices except the sources is non-negative in  $G(\Pi, F)$ . To prove that this assignment of weights makes (21) to simplify to  $-1 \geq 0$  observe that the second sum in (21) is equal to the right hand side of (8) divided by  $B(s)$ . Hence (21) can be written as

$$-Z_s + \sum_{u \in J/\{s\}} \frac{B(u)}{B(s)} \cdot (1 - Z_u) + (Z_s - 1) + \sum_{w \in I/\{s\}} -\frac{B(u)}{B(s)} (1 - Z_u) \geq 0. \quad (22)$$

This completes the first step of the simulation.

*Second step.* The second step is a direct application of Theorem 14: Define the non-negative integers  $b'_u = b_u \cdot B(s)$  and  $c'_w = c_w \cdot B(s)$ . Start at  $\mathcal{H}' = \{I(\ell_u) : u \in J\} \cup \{I(\ell_w) : w \in I\}$ . By Points 4 and 10 in Theorem 14, obtain Frege proofs of  $I(b'_u \cdot \ell_u)$  and  $I(c'_w \cdot \ell_w)$  for each  $u \in J$  and each  $w \in I$ . Now let  $\ell'$  denote the positive linear combination defined as in (21) with  $b_u$  and  $c_w$  replaced by  $b'_u$  and  $c'_w$ , respectively. Recall that  $\ell$  is  $-1 \geq 0$  and hence  $\ell'$  is  $-B(s) \geq 0$ . By Point 3, obtain a Frege proof of  $I(-B(s) \geq 0)$ . By Point 5, obtain a Frege proof of  $I(-1 \geq 0)$ , or equivalently  $I(0 \geq 1)$ . Finally, by Point 11, obtain the Frege proof of 0.

*Third step.* We start the third step by applying the substitution defined by  $Z_u := A_u$  to the refutation of  $\mathcal{H}'$ , where  $A_u$  is again the formula that labels the formula-vertex  $u$ . For each  $v \in I \cup J$ , let  $I(\ell_v)^*$  denote the result of applying this substitution to  $I(\ell_v)$ . To complete the step we need to get polynomial-size Frege proofs of  $I(\ell_v)^*$  from  $\mathcal{H}$ , for each  $v \in I \cup J$ . We do this as a *less direct* application of Theorem 14.

For each formula-vertex  $u \in J$  of a hypothesis formula in  $\mathcal{H}$ , we get a Frege proof of  $I(\ell_u)^*$  from  $A_u$  by applying the substitution  $X := A_u$  to the Frege proof given by Point 6 in Theorem 14. When  $u$  is the formula-vertex of the derived empty formula, we get a Frege proof of  $I(\ell_u)^*$  from  $\bar{0}$  by applying the substitution  $X := 0$  to the Frege proof given by Point 7 in Theorem 14. Since  $\bar{0}$  is the conclusion of an instance of the axiom rule of Frege (namely  $0 \vee \bar{0}$ ), this is a Frege proof of  $I(\ell_u)^*$  from nothing. For every other formula-vertex  $u$ , we get a Frege proof of  $I(\ell_u)^*$  from nothing by applying the substitution  $X := A_u$  to the Frege proof given by Point 8 in Theorem 14.

For each inference-vertex  $w \in I$ , with labelling rule  $R$  and in- and out-neighbours  $N^-$  and  $N^+$ , we proceed as follows. By Point 2 in Theorem 14 and the soundness of  $R$ , first note that  $I(\ell_w)^*$  is a propositional tautology. We claim that, in addition, this tautology is obtained by applying a substitution to another tautology  $T$  that has at most two propositional variables  $X$  and  $Y$ . Concretely,  $T$  will itself be the result of applying a substitution to  $I(\ell_w)$ . We define  $T$  by cases depending on what rule  $R$  is. If  $R$  is the axiom rule and  $N^+ = \{a\}$ , then we take  $T$  to be the result of applying the substitution  $Z_a := X \vee \bar{X}$  to  $I(\ell_w)$ . If  $R$  is the cut rule,  $N^- = \{a, b\}$  and  $N^+ = \{c\}$ , then we take  $T$  to be the result of applying the substitution  $Z_a := Y \vee X$ ,  $Z_b := Y \vee \bar{X}$ , and  $Z_c := Y$  to  $I(\ell_w)$ . If  $R$  is the split rule,  $N^- = \{a\}$  and  $N^+ = \{b, c\}$ , then we take  $T$  to be the result of applying the substitution  $Z_a := Y$ ,  $Z_b := Y \vee X$ , and  $Z_c := Y \vee \bar{X}$  to  $I(\ell_w)$ . By Point 2 in Theorem 14 and the soundness of  $R$ , in all three cases  $T$  is a tautology with at most two propositional variables. By the

completeness of Frege,  $T$  has a constant-size Frege proof. Applying the substitution that turns  $T$  into  $I(\ell_w)^*$  to this proof we get a polynomial-size Frege proof of  $I(\ell_w)^*$  as desired. This completes the third step, and the proof.

## 6 Concluding Remarks

One immediate consequence of Theorem 5 is that there is a polynomial-time algorithm that automates the search for Circular Resolution proofs of bounded width:

**Corollary 15.** *There is an algorithm that, given an integer parameter  $w$  and a set of clauses  $A_1, \dots, A_m$  and  $A$  with  $n$  variables, returns a Circular Resolution proof of width  $w$  of clause  $A$  from  $A_1, \dots, A_m$ , if there is one, and the algorithm runs in time polynomial in  $m$  and  $n^w$ .*

The proof-search algorithm of Corollary 15 relies on linear programming because it relies on our translations to and from Sherali-Adams, whose automating algorithm does rely on linear programming [31]. Based on the fact that the number of clauses of width  $w$  is about  $n^w$ , a direct proof of Corollary 15 is also possible, but as far as we see it still relies on linear programming for finding the flow assignment. It remains as an open problem whether a more combinatorial algorithm exists for the same task.

Another consequence of the equivalence with Sherali-Adams is that Circular Resolution has a length-width relationship in the style of the one for Dag-like Resolution [6]. This follows from Theorem 5 in combination with the size-degree relationship that is known to hold for Sherali-Adams (see [27, 2]). Combining this with the known lower bounds for Sherali-Adams (see [20, 27]), we get the following:

**Corollary 16.** *There are families of 3-CNF formulas  $(F_n)_{n \geq 1}$ , where  $F_n$  has  $O(n)$  variables and  $O(n)$  clauses, such that every Circular Resolution refutation of  $F_n$  has width  $\Omega(n)$  and size  $2^{\Omega(n)}$ .*

It should be noticed that, unlike the well-known observation that tree-like and dag-like width are equivalent measures for Resolution, these are *not* equivalent to width in Circular Resolution. The sparse graph pigeonhole principle from Section 4 illustrates the point. This shows that Circular Resolution proofs of bounded-width proofs cannot be *unfolded* into bounded-width tree-like Resolution proofs.

This observation also explains, perhaps, why our proof that Circular Frege simulates Tree-like Frege goes via a very indirect translation. It also raises one further question (and answer). It is known that Tree-like Bounded-Depth Frege simulates Dag-like Bounded-Depth Frege, at the cost of increasing the depth by one. Could the simulation of Circular Frege by Tree-like Frege be made to preserve bounded depth? The (negative) answer is also provided by the pigeonhole principle which is known to be hard for Bounded-Depth Frege [1, 28, 24], but is easy for Circular Resolution, and hence for Circular Depth-1 Frege.

One last aspect of the equivalence between Circular Resolution and Sherali-Adams concerns the theory of SAT-solving. As is well-known, state-of-the-art SAT-solvers produce Resolution proofs as certificates of unsatisfiability and, as a result, will not be able to handle counting arguments of pigeonhole type. This has motivated the study of so-called *pseudo-Boolean solvers* that handle counting constraints and reasoning through specialized syntax and inference rules. The equivalence between Circular Resolution and Sherali-Adams suggests a completely different approach to incorporate counting capabilities: instead of enhancing the syntax, keep it to clauses but enhance the *proof-shapes*. Whether circular proof-shapes can be handled in a sufficiently effective and efficient way is of course in doubt, but certainly a question worth studying.

It turns out that Circular Resolution has unexpected connections with Dual Rail MaxSAT Resolution [22].

MaxSAT Resolution is a variant of resolution where proofs give upper bounds on the number of clauses of the CNF that can be satisfied simultaneously. At the very least, when the upper bound is less than the number of clauses, MaxSAT resolution provides a refutation of the formula. The Dual Rail encoding is a special encoding of CNF formulas, and Dual Rail MaxSAT Resolution is defined to be MaxSAT resolution applied to the Dual Rail encoding of the input formula. It is argued in [7] that Dual Rail encoding gives strength to the proof system, providing a polynomial refutation of the pigeonhole principle formula. Following the conference version of this paper [4], it was argued in [36] that Circular Resolution polynomially simulates Dual Rail MaxSAT Resolution, in the sense that when the Dual Rail encoding of a CNF formula  $F$  has a MaxSAT Resolution refutation of length  $\ell$  and width  $w$ , then  $F$  has a Circular Resolution refutation of length  $O(\ell w)$ . This is interesting per se and provides yet another proof of Corollary 8. The exact relative strength of Circular Resolution, Dual Rail MaxSAT, and other systems for MaxSAT is studied further in [9, 25].

**Acknowledgments** Both authors were partially funded by European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme, grant agreement ERC-2014-CoG 648276 (AUTAR). First author partially funded by MINECO through TIN2013-48031-C4-1-P (TASSAT2). We acknowledge the work of Jordi Coll who conducted experimental results for finding and visualizing actual circular resolution proofs of small instances of the sparse pigeonhole principle. We thank Moritz Müller for comments and for carefully reading an earlier version of the paper.

## References

- [1] Miklos Ajtai. The complexity of the pigeonhole principle. In *29th Annual Symposium on Foundations of Computer Science*, pages 346–355, Oct 1988.
- [2] A. Atserias and T. Hakoniemi. Size-Degree Trade-offs for Sums-of-Squares and Positivstellensatz Proofs. To appear in *Proceedings of 34th Annual Conference on Computational Complexity (CCC 2019)*. Long version in arXiv:1811.01351 [cs.CC], 2018.
- [3] A. Atserias, M. Lauria, and J. Nordström. Narrow proofs may be maximally long. *ACM Trans. Comput. Log.*, 17(3):19:1–19:30, 2016.
- [4] Albert Atserias and Massimo Lauria. Circular (yet sound) proofs. In Mikolás Janota and Inês Lynce, editors, *Theory and Applications of Satisfiability Testing - SAT 2019 - 22nd International Conference, SAT 2019, Lisbon, Portugal, July 9-12, 2019, Proceedings*, volume 11628 of *Lecture Notes in Computer Science*, pages 1–18. Springer, 2019.
- [5] Yu Hin Au and L. Tunçel. A comprehensive analysis of polyhedral lift-and-project methods. *SIAM J. Discrete Math.*, 30(1):411–451, 2016.
- [6] E. Ben-Sasson and A. Wigderson. Short proofs are narrow - resolution made simple. *J. ACM*, 48(2):149–169, 2001.
- [7] M. L. Bonet, S. Buss, A. Ignatiev, J. Marques-Silva, and A. Morgado. MaxSAT Resolution With the Dual Rail Encoding. In *Proc. 32nd AAAI Conference on Artificial Intelligence*, 2018.
- [8] M. L. Bonet, J. L. Esteban, N. Galesi, and J. Johannsen. On the relative complexity of resolution refinements and cutting planes proof systems. *SIAM J. Comp.*, 30(5), 2000.

- [9] Maria Luisa Bonet and Jordi Levy. Equivalence between systems stronger than resolution. In Luca Pulina and Martina Seidl, editors, *Theory and Applications of Satisfiability Testing - SAT 2020 - 23rd International Conference, Alghero, Italy, July 3-10, 2020, Proceedings*, volume 12178 of *Lecture Notes in Computer Science*, pages 166–181. Springer, 2020.
- [10] J. Brotherston. *Sequent Calculus Proof Systems for Inductive Definitions*. PhD thesis, University of Edinburgh, November 2006.
- [11] J. Brotherston and A. Simpson. Sequent calculi for induction and infinite descent. *Journal of Logic and Computation*, 21(6):1177–1216, 2011.
- [12] Samuel R. Buss. Polynomial size proofs of the propositional pigeonhole principle. *Journal of Symbolic Logic*, 52(4):916–927, 1987.
- [13] Stephen A. Cook and Robert A. Reckhow. The relative efficiency of propositional proof systems. *J. Symb. Log.*, 44(1):36–50, 1979.
- [14] S. S. Dantchev. Rank complexity gap for Lovász-Schrijver and Sherali-Adams proof systems. In *Proc. 39th Annual ACM Symposium on Theory of Computing*, pages 311–317, 2007.
- [15] S. S. Dantchev, B. Martin, and M. N. C. Rhodes. Tight rank lower bounds for the Sherali-Adams proof system. *Theor. Comput. Sci.*, 410(21-23):2054–2063, 2009.
- [16] Anupam Das. On the logical complexity of cyclic arithmetic. *Log. Methods Comput. Sci.*, 16(1), 2020.
- [17] C. Dax, M. Hoffman, and M. Lange. *A Proof System for the Linear Time  $\mu$ -Calculus*, pages 273–284. Springer Berlin Heidelberg, Berlin, Heidelberg, 2006.
- [18] Jérôme Fortier. *Puissance expressive des preuves circulaires. (Expressive Power of Circular Proofs)*. PhD thesis, Aix-Marseille University, Aix-en-Provence, France, 2014.
- [19] Andreas Goerdt. Cutting plane versus frege proof systems. In Egon Börger, Hans Kleine Büning, Michael M. Richter, and Wolfgang Schönfeld, editors, *Computer Science Logic: 4th Workshop, CSL '90 Heidelberg, Germany, October 1–5, 1990 Proceedings*, pages 174–194, Berlin, Heidelberg, 1991. Springer Berlin Heidelberg.
- [20] D. Grigoriev. Linear lower bound on degrees of positivstellensatz calculus proofs for the parity. *Theoretical Computer Science*, 259(1):613 – 622, 2001.
- [21] A. Haken. The intractability of resolution. *Theor. Comp. Sci.*, 39:297 – 308, 1985.
- [22] Alexey Ignatiev, António Morgado, and João Marques-Silva. On tackling the limits of resolution in SAT solving. In *Proc. 20th International Conference on Theory and Applications of Satisfiability Testing - SAT 2017*, pages 164–183, 2017.
- [23] J. Krajíček. *Bounded Arithmetic, Propositional Logic, Complexity Theory*. Cambridge, 1994.
- [24] Jan Krajíček, Pavel Pudlák, and Alan Woods. An exponential lower bound to the size of bounded depth frege proofs of the pigeonhole principle. *Random Structures and Algorithms*, 7(1):15–39, 1995.
- [25] Javier Larrosa and Emma Rollon. Towards a better understanding of (partial weighted) maxsat proof systems. In Luca Pulina and Martina Seidl, editors, *Theory and Applications of Satisfiability Testing -*

- SAT 2020 - 23rd International Conference, Alghero, Italy, July 3-10, 2020, Proceedings*, volume 12178 of *Lecture Notes in Computer Science*, pages 218–232. Springer, 2020.
- [26] D. Niwiński and I. Walukiewicz. Games for the  $\mu$ -calculus. *Theor. Comp. Sci.*, 163(1), 1996.
- [27] T. Pitassi and N. Segerlind. Exponential Lower Bounds and Integrality Gaps for Tree-Like Lovász-Schrijver Procedures. *SIAM J. Comput.*, 41(1):128–159, 2012.
- [28] Toniann Pitassi, Paul Beame, and Russell Impagliazzo. Exponential lower bounds for the pigeonhole principle. *Computational Complexity*, 3:97–140, 1993.
- [29] Alexander Schrijver. *Combinatorial optimization: polyhedra and efficiency*, volume 24. Springer Science & Business Media, 2003.
- [30] Daniyar S. Shamkanov. Non-well-founded derivations in the gödel-löb provability logic. *Rev. Symb. Log.*, 13(4):776–796, 2020. Early draft as <http://arxiv.org/abs/1401.4002> in 2014.
- [31] H. D. Sherali and W. P. Adams. A hierarchy of relaxations between the continuous and convex hull representations for zero-one programming problems. *SIAM J. Disc. Math.*, 3(3):411–430, 1990.
- [32] J. Shoesmith and T. J. Smiley. *Multiple-Conclusion Logic*. Cambridge, 1978.
- [33] Alex Simpson. Cyclic arithmetic is equivalent to peano arithmetic. In Javier Esparza and Andrzej S. Murawski, editors, *Foundations of Software Science and Computation Structures - 20th International Conference, FOSSACS 2017, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2017, Uppsala, Sweden, April 22-29, 2017, Proceedings*, volume 10203 of *Lecture Notes in Computer Science*, pages 283–300, 2017.
- [34] T. Studer. On the proof theory of the modal mu-calculus. *Studia Logica*, 89(3), 2008.
- [35] W. W. Tait. Normal derivability in classical logic. In J. Barwise, editor, *The Syntax and Semantics of Infinitary Languages*, volume 72 of *Lecture Notes in Mathematics*, pages 204–236. Springer-Verlag, 1968.
- [36] M. Vinyals. Personal communication, 2018.