

ON THE MAXIMAL PART IN UNREFINABLE PARTITIONS OF TRIANGULAR NUMBERS

RICCARDO ARAGONA¹, LORENZO CAMPIONI¹, ROBERTO CIVINO¹ AND MASSIMO LAURIA²

ABSTRACT. A partition into distinct parts is refinable if one of its parts a can be replaced by two different integers which do not belong to the partition and whose sum is a , and it is unrefinable otherwise. Clearly, the condition of being unrefinable imposes on the partition a non-trivial limitation on the size of the largest part and on the possible distributions of the parts. We prove a $O(n^{1/2})$ -upper bound for the largest part in an unrefinable partition of n , and we call maximal those which reach the bound. We show a complete classification of maximal unrefinable partitions for triangular numbers, proving that if n is even there exists only one maximal unrefinable partition of $n(n+1)/2$, and that if n is odd the number of such partitions equals the number of partitions of $\lceil n/2 \rceil$ into distinct parts. In the second case, an explicit bijection is provided.

1. INTRODUCTION

Integer partitions into distinct parts may appear in several areas of mathematics, sometimes unexpectedly. For example, they have been recently shown to be linked to the set of generators of groups in a group-theoretical problem related to cryptography [ACGS19, ACGS21a]. In particular, Aragona et al. showed that the generators of a given group are linked to partitions into distinct parts which satisfy a condition of *non-refinability* [ACGS21b] together with a condition on the minimal excludant. This motivates us to investigate some combinatorial aspects of *unrefinable partitions*, i.e. those in which no part can be written as the sum of two different integers which do not belong to the partition, which to our knowledge have not been investigated so far (cf. the On-Line Encyclopedia of Integer Sequences for the first values [OEI, <https://oeis.org/A179009>]).

Computational results suggest that the maximal part in an unrefinable partition of n is approximatively \sqrt{n} . In this paper, we first prove a matching upper bound for the maximal part and then we define *maximal unrefinable partitions* as those which reach the bound. As a main contribution, we provide a complete classification of maximal unrefinable partitions for triangular numbers. We constructively prove, denoting by T_n the n -th triangular number, that for even n there exists exactly one maximal unrefinable partition of T_n . For odd n , we obtain a lower bound for the minimal excludant for the maximal unrefinable partitions of T_n , defined to be the least integers which is not a part [FP15] and which has been investigated also recently by other authors [AN19, HSS22]. The knowledge of a bound on the minimal

¹ DISIM - UNIVERSITÀ DEGLI STUDI DELL'AQUILA, ITALY

² DIPARTIMENTO DI SCIENZE STATISTICHE - SAPIENZA, UNIVERSITÀ DI ROMA, ITALY

E-mail addresses: riccardo.aragona@univaq.it, roberto.civino@univaq.it, lorenzo.campioni1@graduate.univaq.it, massimo.lauria@uniroma1.it.

2010 *Mathematics Subject Classification.* 11P81, 05A17, 05A19.

Key words and phrases. Unrefinable partitions, partitions into distinct parts, triangular numbers, minimal excludant, bijective proof.

R. Aragona and R. Civino are members of INdAM-GNSAGA (Italy). R. Civino is partially funded by the Centre of excellence ExEMERGE at the University of L'Aquila.

excludant, among other considerations, allows us to show an explicit bijection between the set of the maximal unrefinable partitions of T_n and the set of partitions of $\lceil n/2 \rceil$ into distinct parts in the classic sense [And76].

The remainder of the paper is organized as follows: in Sec. 2 we introduce the notation and define unrefinable partitions. In Sec. 3 we prove two upper bounds for the maximal part in an unrefinable partition of n , distinguishing the case when n is a triangular number and when is not. The classification theorem, i.e. Theorem 4.1, is proved in Sec. 4, which also contains the result on triangular numbers of an even number. The odd case is developed in Sec 5, which concludes the paper. In particular, we show in Theorem 5.11 a bijective proof that number of maximal unrefinable partitions of T_n equals the number of partitions of $\lceil n/2 \rceil$ into distinct parts.

2. PRELIMINARIES

Let $N \in \mathbb{N}$. A *partition* of N is a finite sequence $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t)$ of positive integers such that $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_t$ and $\sum_{i=1}^t \lambda_i = N$. When λ is partition of N we write $\lambda \vdash N$. Each λ_i is called a *part* of the partition λ and we call λ_t its *maximal part*. We denote by $(\lambda_1, \lambda_2, \dots, \lambda_{i-1}, \widehat{\lambda}_i, \lambda_{i+1}, \dots, \lambda_t)$ the partition $(\lambda_1, \lambda_2, \dots, \lambda_{i-1}, \lambda_{i+1}, \dots, \lambda_t)$ where the part λ_i is removed.

The partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t)$ is a *partition into distinct parts* if $\lambda_1 < \lambda_2 < \dots < \lambda_t$ and $t \geq 2$, i.e. if each part appears exactly once. The set \mathbb{D}_N denotes the set of all the partitions of N into distinct parts. If $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t) \in \mathbb{D}_N$, the integers belonging to

$$\mathcal{M}_\lambda \stackrel{\text{def}}{=} \{1, 2, \dots, \lambda_t\} \setminus \{\lambda_1, \lambda_2, \dots, \lambda_t\}$$

are called the *missing parts* of λ , and are denoted by $\mu_1 < \mu_2 < \dots < \mu_m$, for some $m \geq 0$. The least integer which is not a part of λ , i.e. μ_1 , is the *minimal excludant* of λ [FP15]. We denote this by writing $\mu_1 = \text{mex}(\lambda)$, taking $\text{mex}(\lambda) = 0$ when $\mathcal{M}_\lambda = \emptyset$ as it is customary in literature.

Definition 2.1. Let $N \in \mathbb{N}$. Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t)$ be a partition of N into distinct parts and let $\mu_1 < \mu_2 < \dots < \mu_m$ be its missing parts. The partition λ is *refinable* if there exist $1 \leq \ell \leq t$ and $1 \leq i < j \leq m$ such that $\mu_i + \mu_j = \lambda_\ell$, and *unrefinable* otherwise. The set of unrefinable partitions is denoted by \mathbb{U} , and by \mathbb{U}_N we denote those whose sum of the parts is N .

Definition 2.2. Let $n \in \mathbb{N}$. We denote by T_n the *n-th triangular number*, i.e.

$$T_n \stackrel{\text{def}}{=} \sum_{i=1}^n i = \frac{n(n+1)}{2}.$$

The *complete partition* $\pi_n \stackrel{\text{def}}{=} (1, 2, \dots, n)$ is the partition of T_n with no missing parts.

Notice that every complete partition is unrefinable. The same holds, by definition, for partitions with a single missing part. In particular, if $N = T_n$ for some n , then π_n is an unrefinable partition of N . Otherwise, if n is the least integer such that $N < T(n)$, then

$$\pi_{n,d} \stackrel{\text{def}}{=} (1, 2, \dots, d-1, \widehat{d}, d+1, \dots, n) \quad (1)$$

is an unrefinable partition of N , where $d = T_n - N$. In general, the admissible number of missing parts in an unrefinable partition is bounded as in the following result.

Lemma 2.3. *Let $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t)$ be unrefinable and let $\mu_1 < \mu_2 < \dots < \mu_m$ be the missing parts. Then the number of missing parts m is bounded by*

$$m \leq \left\lfloor \frac{\lambda_t}{2} \right\rfloor. \quad (2)$$

Proof. Let us start by observing that $\lambda_t - \mu_i \in \lambda$ for $1 \leq i \leq m$, otherwise from $\lambda_t - \mu_i, \mu_i \in \mathcal{M}_\lambda$ we obtain $(\lambda_t - \mu_i) + \mu_i = \lambda_t \in \lambda$ and thus λ is refinable. We prove the claim considering the complete partition π_{λ_t} and removing from this the maximum number of parts different from λ_t . For the previous observation, each candidate part μ_i to be removed has a counterpart $\lambda_t - \mu_i$ in the partition. The bound of Eq. (2) depends on the fact that this process can be repeated no more than $\lfloor \lambda_t/2 \rfloor$ times. \square

As already anticipated, our focus is on the maximal part in a partition as in Definition 2.1. In the next section, using Lemma 2.3, we show that the maximal part in an unrefinable partition of n is $O(n^{1/2})$.

3. UPPER BOUNDS ON THE MAXIMAL PART

It is easy to check that the complete partitions $\pi_1, \pi_2, \dots, \pi_5$ are the only unrefinable partitions for the triangular numbers T_1, T_2, \dots, T_5 respectively. In the general case of T_n for $n \geq 6$, this is not true. For example, the partition $(1, 2, 3, 7, 8) \vdash 21 = T_6$ is unrefinable. As a more complex example, in case of T_9 we can calculate that

$$\begin{aligned} (1, 2, 3, 4, 5, 6, 7, 8, 9) & (1, 2, 3, 5, 6, 7, 10, 11) \\ (1, 2, 3, 4, 6, 8, 10, 11) & (1, 2, 3, 4, 5, 9, 10, 11) \\ (1, 2, 3, 4, 6, 7, 10, 12) & (1, 2, 3, 4, 5, 8, 10, 12) \\ (1, 2, 3, 4, 5, 7, 11, 12) & (1, 2, 3, 4, 5, 7, 10, 13) \\ (1, 2, 3, 4, 5, 6, 11, 13) & (1, 2, 3, 4, 5, 6, 10, 14) \\ (1, 2, 4, 5, 8, 11, 14) & \end{aligned}$$

are all the unrefinable partitions of $45 = T_9$.

It is clear that the property of being unrefinable imposes on one side an upper limitation on the size of the largest part which is admissible in the partition, and on the other a lower limitation on the minimal excludant. We address in this section the natural question of determining what is the maximal part in an unrefinable partition of N . In the case where N is a triangular number the following result provides an answer. The notation introduced in the proof will be used throughout all the paper.

Proposition 3.1. *Let $n \in \mathbb{N}$ and $N = T_n$. For every $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t)$ unrefinable partition of N we have*

$$n \leq \lambda_t \leq 2n - 4. \quad (3)$$

Equivalently,

$$\frac{\sqrt{1+8N}-1}{2} \leq \lambda_t \leq \sqrt{1+8N}-5.$$

Proof. Let us start considering the complete partition $\pi_n \vdash N$. Other unrefinable partitions of N are obtained from π_n by removing some parts smaller or equal than n and replacing them with parts larger than n . Hence, the lower bound for the maximal part in any partition

of N is n , obtained when no part is removed. Since $N = n(n+1)/2$, then n is the positive solution of $n^2 + n - 2N = 0$ and so we have

$$\lambda_t \geq \frac{\sqrt{1+8N}-1}{2}.$$

Let $h, j \in \mathbb{N}$ and let us denote by $1 \leq a_1 < a_2 < \dots < a_h \leq n$ the candidate parts to be removed from π_n to obtain a new unrefinable partition of N , and by $n+1 \leq \alpha_1 < \alpha_2 < \dots < \alpha_j$ the corresponding replacements. Since $\sum a_i = \sum \alpha_i$ we have $h > j$. Moreover $j > 1$, otherwise from $\sum a_i = \alpha_1$ the obtained partition is refinable. Hence we obtain

$$h \geq 3, j \geq 2, \text{ and } h > j.$$

There are h missing parts in the interval $\{1, 2, \dots, n\}$ and exactly j parts appear in the interval $\{n+1, n+2, \dots, \alpha_j\}$. Therefore the number of missing parts of λ is

$$m = h + \alpha_j - n - j.$$

To prove $\alpha_j \leq 2n - 4$ we consider the cases where α_j is either equal to $2n - 3$, equal to $2n - 2$, or strictly larger than $2n - 2$. We derive a contradiction in each case. Let us observe that

$$\sum_{i=1}^h a_i \leq n + (n-1) + \dots + (n - (h-1)) = hn - \frac{(h-1)h}{2}.$$

In the case $\alpha_j = 2n - 3$ we obtain $m = h + n - 3 - j$. By Lemma 2.3, we have $m \leq \lfloor \alpha_j/2 \rfloor = n - 2$, hence $h \leq j + 1$, and so $h = j + 1$. Notice that

$$\alpha_1 + \dots + \alpha_j > (j-1)n + 2n - 3 = (j+1)n - 3 = hn - 3.$$

Therefore, since $\sum a_i = \sum \alpha_i$, we have

$$3 > \frac{(h-1)h}{2},$$

which is satisfied if $h < 3$, a contradiction.

In the case $\alpha_j = 2n - 2$ we obtain $m = h + n - 2 - j$. By Lemma 2.3, we have $m \leq \lfloor \alpha_j/2 \rfloor = n - 1$, hence $h \leq j + 1$, and so again $h = j + 1$. Notice that

$$\alpha_1 + \dots + \alpha_j > (j-1)n + 2n - 2 = (j+1)n - 2 = hn - 2.$$

Therefore, since $\sum \alpha_i = \sum a_i$, we have

$$2 > \frac{(h-1)h}{2},$$

which is satisfied if $h < (1 + \sqrt{17})/2 < 3$, a contradiction.

To conclude we consider the last case $\alpha_j > 2n - 2$. We have $n - 1 < \alpha_j/2$ and so

$$\alpha_j - (n-1) > \frac{\alpha_j}{2} \geq \left\lfloor \frac{\alpha_j}{2} \right\rfloor.$$

Hence, since $h \geq j + 1$, we have

$$m = h + \alpha_j - n - j > \left\lfloor \frac{\alpha_j}{2} \right\rfloor + h - j - 1 \geq \left\lfloor \frac{\alpha_j}{2} \right\rfloor,$$

which contradicts Lemma 2.3. □

Notice that the upper bound of Eq. 3 is tight. Indeed, let us define the following partition:

$$\widetilde{\pi}_n \stackrel{\text{def}}{=} (1, 2, \dots, n-3, n+1, 2n-4). \quad (4)$$

It is easy to notice that $\widetilde{\pi}_n \vdash N$ and that $\widetilde{\pi}_n$ is unrefinable, since its least missing parts are $n-2$ and $n-1$, and $2n-4 < (n-2) + (n-1)$. In the notation of the proof of Proposition 3.1, $\widetilde{\pi}_n$ is obtained in the case $h = 3$ and $j = h - 1 = 2$.

The counterpart of Proposition 3.1 in the case of non-triangular numbers is obtained in a similar way.

Proposition 3.2. *Let $N \in \mathbb{N}$ be such that $T_{n-1} < N < T_n$ for some $n \in \mathbb{N}$. For every $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t)$ unrefinable partition of N we have*

$$n \leq \lambda_t \leq 2n - 2. \quad (5)$$

Equivalently,

$$\frac{\sqrt{1 + 8(N + d)} - 1}{2} \leq \lambda_t \leq \sqrt{1 + 8(N + d)} - 3,$$

where $d = T_n - N$.

Proof. Let us start considering d and the partition $\pi_{n,d} \vdash N$ as in Eq. (1). Other partitions of N are obtained from $\pi_{n,d}$ by removing some parts smaller or equal than n which are replaced by d and other parts larger than n or only by other parts larger than n . Proceeding as in the proof of Proposition 3.1, let $h, j \in \mathbb{N}$ and let us denote by $1 \leq a_1 < a_2 < \dots < a_h \leq n$ the candidate parts to be removed from $\pi_{n,d}$ to obtain a new partition of N , and by $\alpha_1 < \alpha_2 < \dots < \alpha_j$ the corresponding replacements. Since $\sum a_i = \sum \alpha_i$ we have $h \geq j > 1$, and we may obtain $h = j$ only if $\alpha_1 = d$. For this reason, we need do consider the two cases separately.

Let us assume $\alpha_i > n$, for every $1 \leq i \leq j$. Reasoning as in the proof of Proposition 3.1 we can count $m = (h + 1) + \alpha_j - n - j$. On the other hand, if $\alpha_1 = d$ and $\alpha_i > n$ for every $2 \leq i \leq j$, then we obtain just h missing parts in the interval $\{1, 2, \dots, n\}$ and exactly $j - 1$ parts appear in the interval $\{n + 1, n + 2, \dots, \alpha_j\}$, therefore we obtain the same formula for number of missing parts $m = h + \alpha_j - n - (j - 1)$. By Lemma 2.3 we obtain

$$h + \left\lceil \frac{\alpha_j}{2} \right\rceil - n - j + 1 \leq 0. \quad (6)$$

If $\alpha_j > 2n - 2$, then $\lceil \alpha_j / 2 \rceil \geq n$ and from Eq. (6) we obtain $h \leq j - 1$, a contradiction. \square

Remark 1. Notice that the bound of Eq. (5) is reached by the partition $(1, 2, \dots, n - 2, 2n - 2) \vdash T_n - 1$ constructed from $\pi_{n,1} = (2, 3, \dots, n)$.

We now introduce maximal unrefinable partitions, the main subject of this work, as those partitions $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t) \in \widetilde{\mathcal{U}}_N$ whose λ_t is maximal.

Definition 3.3. Let $N \in \mathbb{N}$. An unrefinable partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t)$ of N is called *maximal* if

$$\lambda_t = \max_{(\lambda'_1, \lambda'_2, \dots, \lambda'_t) \in \mathcal{U}_N} \lambda'_t$$

We denote by $\widetilde{\mathcal{U}}_N$ the set of the maximal unrefinable partitions of N .

In the case of triangular numbers $N = T_n$ for some $n \geq 6$, by virtue of Proposition 3.1, we have that $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t) \in \widetilde{\mathcal{U}}_N$ is maximal if and only if $\lambda_t = 2n - 4$.

Remark 2. As already observed in the proof of Proposition 3.1, for each $\lambda \in \mathcal{U}$ there exist $1 < j < h$, $1 \leq a_1 < a_2 < \dots < a_h \leq n$ and $\alpha_1, \alpha_2, \dots, \alpha_j \geq n + 1$ such that λ is obtained from π_n by removing the parts a_i s which are replaced by the parts α_i s. Consequently, $\#\widetilde{\mathcal{U}}_N$ coincides with the number of choices which lead to partitions meeting the mentioned conditions and, in addition, the condition $\lambda_t = 2n - 4$. In the remainder of the work, when $\lambda \in \mathcal{U}$ we will refer to a_i s, α_i s, j and h as intended here.

4. CLASSIFICATION OF MAXIMAL UNREFINABLE PARTITIONS OF TRIANGULAR NUMBERS

We are now ready to prove our first main contribution. Using arguments similar to those of the proofs in the previous section, we classify maximal unrefinable partitions for triangular numbers.

Theorem 4.1. *Let $n \in \mathbb{N}$, $n \geq 6$, and $N = T_n$. Then*

- (1) *if n is even, then $\widetilde{\mathcal{U}}_N = \{\widetilde{\pi}_n\}$;*
- (2) *if n is odd, then $\widetilde{\pi}_n \in \widetilde{\mathcal{U}}_N$ and the other partitions $\lambda \in \widetilde{\mathcal{U}}_N$, $\lambda \neq \widetilde{\pi}_n$, are such that $j = h - 2$ and the following conditions hold:*
 - (i) *the removed parts a_1, \dots, a_{h-3} are replaced by $2n - 4 - a_1, 2n - 4 - a_2, \dots, 2n - 4 - a_{h-3}$, and*
 - (ii) *the triple (a_{h-2}, a_{h-1}, a_h) is one of the following*
 $(n - 4, n - 3, n - 2), (n - 4, n - 2, n - 1), (n - 3, n - 2, n), (n - 2, n - 1, n)$.

Proof. Let $\lambda \in \widetilde{\mathcal{U}}_N$ and let a_1, a_2, \dots, a_h and $\alpha_1, \alpha_2, \dots, \alpha_j = 2n - 4$ as in Remark 2. We already know that $h \geq 3$. From the hypotheses on λ we have that

$$m = h + \alpha_j - n - j = h + n - 4 - j.$$

By Lemma 2.3 we have $h - j \leq 2$, and, since $h > j$, we obtain $j \in \{h - 1, h - 2\}$. Notice that if $a \in \{a_1, \dots, a_h\}$ is such that $a < n - 4$, then $\alpha \stackrel{\text{def}}{=} 2n - 4 - a$ must belong to $\{\alpha_1, \dots, \alpha_{j-1}\}$, otherwise $\alpha + a = 2n - 4 = \alpha_j \in \lambda$, and so λ is refinable. Then each removed part a_i such that $a_i < n - 4$ is in one-to-one correspondence to its *replacement* which, for the sake of simplicity, we will denote from now on by α_i . On the other side, for the same symmetry argument, each part in the interval $\{n - 4, \dots, n\}$ has no replacement. In such an interval we may choose at most 5 parts. However, we are not allowed to remove, at the same time, parts from one of the pairs $(n - 4, n)$ and $(n - 3, n - 1)$ without contradicting the unrefinability of λ . Analogously, we are not allowed to remove more than three parts. Moreover, we cannot choose to pick only one part to be removed in that interval, otherwise we would obtain $h - 1$ replacements but at most $j - 1$ are allowed, and $h > j$.

We are left to consider the cases of two or three parts to be removed in the interval $\{n - 4, \dots, n\}$, both with the assumptions $j = h - 1$ or $j = h - 2$. In particular, we will show that in both setting of j , there exists no maximal partition with two removed parts in the selected interval. Moreover, in the case $j = h - 1$ and three removed parts, we show that the only admissible partition is $\widetilde{\pi}_n$. Finally, partitions from the case $j = h - 2$ and three removed parts are only possible for odd n as claimed in (2). Let us address each of the four cases separately.

Let us suppose $j = h - 1$ and $1 \leq a_1 < a_2 < \dots < a_{h-2} \leq n - 5$ and $n - 4 \leq a_{h-1} < a_h \leq n$. For each $1 \leq i \leq j - 1 = h - 2$ we have $\alpha_i = \alpha_j - a_i$. We will now show that this configuration leads to a contradiction. To do this, we estimate $\sum a_i$ and $\sum \alpha_i$ from above and from below, respectively. This is clearly accomplished by noticing that $\alpha_{h-2} \geq n + 1, \alpha_{h-3} \geq n + 2, \dots, \alpha_1 \geq n + h - 2$ and $a_h \leq n, a_{h-1} \leq n - 1$, obtaining $a_{h-2} \leq \alpha_j - \alpha_{h-2} \leq n - 5, a_{h-3} \leq n - 6, \dots, a_1 \leq n - h - 2$. Hence

$$\sum_{i=1}^h a_i \leq hn - \sum_{i=1}^{h+2} i + 2 + 3 + 4 = hn - \frac{h^2 + 5h + 6}{2} + 9, \text{ and}$$

$$\sum_{i=1}^j \alpha_i \geq hn + \sum_{i=1}^{h-2} i - 4 = hn + \frac{h^2 - 3h + 2}{2} - 4.$$

For $\sum a_i = \sum \alpha_i$ we obtain an inequality which is satisfied for $h < 3$, which is a contradiction.

In the second case, i.e. $j = h - 1$, and $1 \leq a_1 < a_2 < \dots < a_{h-3} \leq n - 5$, $n - 4 \leq a_{h-2} < a_{h-1} < a_h \leq n$, we have $\alpha_i = \alpha_j - a_i$ for every $1 \leq i \leq h - 3 = j - 2$. Notice that, in this case, the part $\alpha_{j-1} = \alpha_{h-2}$ is not determined by one of the a_i s. Proceeding as before, since $\alpha_{h-3} \geq n + 1$, $\alpha_{h-4} \geq n + 2, \dots, \alpha_1 \geq n + h - 3$, $\alpha_{h-2} \geq n + h - 2$ and $a_h \leq n, a_{h-1} \leq n - 1, a_{h-2} \leq n - 2$, we determine $a_{h-3} \leq \alpha_j - \alpha_{h-3} \leq n - 5, a_{h-4} \leq n - 6, \dots, a_1 \leq n - h - 1$ and we obtain the bounds

$$\sum a_i \leq hn - \sum_{i=1}^{h+1} i + 3 + 4 = hn - \frac{h^2 + 3h + 2}{2} + 7, \text{ and}$$

$$\sum \alpha_i \geq hn + \sum_{i=1}^{h-2} i - 4 = hn + \frac{h^2 - 3h + 2}{2} - 4.$$

From $\sum a_i = \sum \alpha_i$ we obtain an inequality which is satisfied only for $h = 3$, which corresponds to the partition (cf. also Eq. (4))

$$(1, 2, \dots, n - 3, n + 1, 2n - 4) = \widetilde{\pi}_n \in \widetilde{\mathcal{U}}_N.$$

The third case $j = h - 2$ with two removed parts is immediately contradictory, since the parts a_1, a_2, \dots, a_{h-2} determine $h - 2 = j$ replacements but at most $j - 1$ are possible.

The last case to be considered is the one where $j = h - 2$ and the three largest parts a_i s are chosen in the interval $\{n - 4, \dots, n\}$. As already observed, since λ is unrefinable, the only possible choices are

$$(a_{h-2}, a_{h-1}, a_h) \in \{(n - 4, n - 3, n - 2), (n - 4, n - 2, n - 1), (n - 3, n - 2, n), (n - 2, n - 1, n)\},$$

which means $a_{h-2} + a_{h-1} + a_h \in \{3n - 9, 3n - 7, 3n - 5, 3n - 3\}$. From $\sum a_i = \sum \alpha_i$ we obtain

$$a_1 + a_2 + \dots + a_h = (\alpha_{h-2} - a_1) + (\alpha_{h-2} - a_2) + \dots + (\alpha_{h-2} - \alpha_{h-3}) + \alpha_{h-2}$$

and so, since $\alpha_{h-2} = \alpha_j = 2n - 4$,

$$2(a_1 + a_2 + \dots + a_{h-3}) + (a_{h-2} + a_{h-1} + a_h) = (h - 2)(2n - 4). \quad (7)$$

Since the right side of Eq. (7) is even and $a_{h-2} + a_{h-1} + a_h$ is even only if n is odd, then Eq. (7) can be satisfied only in the case when n is odd. This proves (2) when n is odd and that the partition $\widetilde{\pi}_n$ of Eq. (4) is the only maximal unrefinable partition of T_n when n is even, i.e. (1). \square

From Theorem 4.1 we obtain that the description of maximal unrefinable partition for the triangular number of an even number is completed. The odd case is addressed in the following section.

Corollary 4.2. *Let $k \in \mathbb{N}$ and $N = T_{2k}$. Then $\#\widetilde{\mathcal{U}}_N = 1$.*

5. ODD TRIANGULAR NUMBERS

Throughout all this last section, N will denote the triangular number of an odd number. More precisely, let $n = 2k - 1 \in \mathbb{N}$ be such that $N = T_n$.

From Theorem 4.1 we have that the set of maximal unrefinable partitions of triangular numbers of odd numbers can be partitioned in the following way

$$\{\widetilde{\pi}_n \mid n \text{ odd}\} \dot{\cup} \mathcal{A} \dot{\cup} \mathcal{B} \dot{\cup} \mathcal{C} \dot{\cup} \mathcal{D},$$

where

$$\mathcal{A} \stackrel{\text{def}}{=} \bigcup_{h \geq 4} \mathcal{A}_h, \quad \mathcal{B} \stackrel{\text{def}}{=} \bigcup_{h \geq 4} \mathcal{B}_h, \quad \mathcal{C} \stackrel{\text{def}}{=} \bigcup_{h \geq 4} \mathcal{C}_h, \quad \mathcal{D} \stackrel{\text{def}}{=} \bigcup_{h \geq 4} \mathcal{D}_h$$

and

$$\begin{aligned} \mathcal{A}_h &\stackrel{\text{def}}{=} \bigcup_{n \text{ odd}} \{\lambda \mid \lambda \in \widetilde{\mathcal{U}}_{T_n}, (a_{h-2}, a_{h-1}, a_h) = (n-4, n-3, n-2)\}, \\ \mathcal{B}_h &\stackrel{\text{def}}{=} \bigcup_{n \text{ odd}} \{\lambda \mid \lambda \in \widetilde{\mathcal{U}}_{T_n}, (a_{h-2}, a_{h-1}, a_h) = (n-4, n-2, n-1)\}, \\ \mathcal{C}_h &\stackrel{\text{def}}{=} \bigcup_{n \text{ odd}} \{\lambda \mid \lambda \in \widetilde{\mathcal{U}}_{T_n}, (a_{h-2}, a_{h-1}, a_h) = (n-3, n-2, n)\}, \\ \mathcal{D}_h &\stackrel{\text{def}}{=} \bigcup_{n \text{ odd}} \{\lambda \mid \lambda \in \widetilde{\mathcal{U}}_{T_n}, (a_{h-2}, a_{h-1}, a_h) = (n-2, n-1, n)\}. \end{aligned}$$

Each set $\mathcal{A}_h, \mathcal{B}_h, \mathcal{C}_h, \mathcal{D}_h$ is called a *class* of maximal unrefinable partitions. If $\lambda \in \mathcal{A}_h$ (resp. $\mathcal{B}_h, \mathcal{C}_h$ or \mathcal{D}_h) for some h we say that λ is a *partition of class* \mathcal{A}_h (resp. $\mathcal{B}_h, \mathcal{C}_h$ or \mathcal{D}_h).

The following consideration is a trivial but important consequence of Theorem 4.1.

Corollary 5.1. *Let $n \in \mathbb{N}$ and $N = T_n$. If $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t) \in \widetilde{\mathcal{U}}_N$, then $\lambda_i \neq n-2$ for every $1 \leq i \leq t$.*

Remark 3 (Anti-symmetric property). From Theorem 4.1(2) and Corollary 5.1 we derive that every partition $\lambda \in \widetilde{\mathcal{U}}_{T_n}, \lambda \neq \widetilde{\pi}_n$, is *anti-symmetric* with respect to $n-2$, i.e. for $1 \leq a < 2n-4$ we have

$$a \notin \lambda \iff 2n-4-a \in \lambda,$$

provided that $a \neq n-2$.

Example 5.2. Let us fix $n = 13$. In Tab. 1 we have displayed the three different partitions of $\widetilde{\mathcal{U}}_{T_{13}} \setminus \{\widetilde{\pi}_{13}\}$, where a black dot means that the corresponding integer is a part and the white dot means otherwise. Disregarding the last part which is fixed to be $2n-4$ due to the maximality constraint, the anti-symmetric property with respect to $n-2$ can be appreciated. Notice also that $\min_{\lambda \in \widetilde{\mathcal{U}}_{T_{13}}} \text{mex}(\lambda) = 5 = (n-3)/2$ and that $(n-2) - 5 + 1 = 7 = \lceil n/2 \rceil$.

										n-2											λ _t	
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	
•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•

TABLE 1. The anti-symmetric property shown on the partitions $\lambda \in \widetilde{\mathcal{U}}_{T_{13}}, \lambda \neq \widetilde{\pi}_{13}$.

Example 5.3. As another significative example, we show in Tab 2 all the partitions in $\widetilde{\mathcal{U}}_{T_{27}} = \widetilde{\mathcal{U}}_{378}$, classified according to the description of Theorem 4.1. It is important to notice that, when $h \geq 5$, partitions in the same class may appear with different multiplicities. Here all the parts λ_i s of the partitions are listed, divided in three *areas* $1 \leq \lambda_i \leq n-5$, $n-4 \leq \lambda_i \leq n$ and $n+1 \leq \lambda_i \leq 2n-4$ naturally induced by Theorem 4.1. Notice again that we have $\min_{\lambda \in \widetilde{\mathcal{U}}_{T_{27}}} \text{mex}(\lambda) = 12 = (n-3)/2$.

$1 \leq \lambda_i < 22$	$23 \leq \lambda_i < 27$	$28 \leq \lambda_i \leq 50$	class
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22	23 24	28 50	$\widetilde{\pi}_{27}$
1 2 3 4 5 6 7 8 9 10 11 12 13 15 16 17 18 19 20 21 22	26 27	36 50	\mathcal{A}_4
1 2 3 4 5 6 7 8 9 10 11 12 14 15 16 17 18 19 20 21 22	24 27	37 50	\mathcal{B}_4
1 2 3 4 5 6 7 8 9 10 11 13 14 15 16 17 18 19 20 21 22	23 26	38 50	\mathcal{C}_4
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 21 22	26 27	30 31 50	\mathcal{A}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 19 20 22	26 27	29 32 50	\mathcal{A}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 18 19 20 21	26 27	28 33 50	\mathcal{A}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 19 21 22	24 27	30 32 50	\mathcal{B}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 18 19 20 22	24 27	29 33 50	\mathcal{B}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 17 18 19 20 21	24 27	28 34 50	\mathcal{B}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 20 21 22	23 26	31 32 50	\mathcal{C}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 18 19 21 22	23 26	30 33 50	\mathcal{C}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 17 18 19 20 22	23 26	29 34 50	\mathcal{C}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 16 17 18 19 20 21	23 26	28 35 50	\mathcal{C}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 18 20 21 22	23 24	31 33 50	\mathcal{D}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 17 18 19 21 22	23 24	30 34 50	\mathcal{D}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 16 17 18 19 20 22	23 24	29 35 50	\mathcal{D}_5
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19	24 27	28 29 30 50	\mathcal{B}_6
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 20	23 26	28 29 31 50	\mathcal{C}_6
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 21	23 24	28 30 31 50	\mathcal{D}_6
1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 19 20	23 24	28 29 32 50	\mathcal{D}_6

TABLE 2. Maximal unrefinable partitions of $378 = T_{27}$ and the corresponding classes.

5.1. Bounds for h and the minimal excludant. It is natural to wonder, recalling that in general $h \geq 4$, what is an upper bound for h in a maximal unrefinable partition $\lambda \in \widetilde{\mathcal{U}}_{T_{2k-1}}$. The answer to this question is provided in Proposition 5.6, from which we also derive the result on the lower bound for the minimal excludant in maximal unrefinable partitions (cf. Corollary 5.9). Let us address before the two extremal cases $h = 4$ and $h = 5$.

Proposition 5.4. *Let $n \geq 7$ be odd. We have:*

- (1) $\widetilde{\mathcal{U}}_{T_n} \cap \mathcal{D}_4 = \emptyset$,
- (2) $\widetilde{\mathcal{U}}_{T_7} \cap \mathcal{C}_4 = \emptyset$ and if $n \geq 9$, then $\#(\widetilde{\mathcal{U}}_{T_n} \cap \mathcal{C}_4) = 1$,
- (3) $\widetilde{\mathcal{U}}_{T_7} \cap \mathcal{B}_4 = \widetilde{\mathcal{U}}_{T_9} \cap \mathcal{B}_4 = \emptyset$ and if $n \geq 11$, then $\#(\widetilde{\mathcal{U}}_{T_n} \cap \mathcal{B}_4) = 1$,
- (4) $\widetilde{\mathcal{U}}_{T_7} \cap \mathcal{A}_4 = \widetilde{\mathcal{U}}_{T_9} \cap \mathcal{A}_4 = \emptyset$ and if $n \geq 11$, then $\#(\widetilde{\mathcal{U}}_{T_n} \cap \mathcal{A}_4) = 1$.

Proof. Let $\lambda \in \widetilde{\mathcal{U}}_{T_n}$ be obtained by removing the integers a_1, a_2, a_3, a_4 and adding the replacements α_1 and $\alpha_2 = 2n - 4$, which need to satisfy the following conditions:

- (i) $1 \leq a_1 \leq n - 5$,
- (ii) $n - 4 \leq a_2 < a_3 < a_4 \leq n$,
- (iii) $n + 1 \leq \alpha_1 = 2n - 4 - a_1$,
- (iv) $a_1 + a_2 + a_3 + a_4 = \alpha_1 + \alpha_2$.

By contradiction, let us assume that $\lambda \in \mathcal{D}_4$, and so $a_2 = n - 2, a_3 = n - 1$ and $a_4 = n$. From Eq. (iv) we obtain

$$a_1 = \frac{n-5}{2} \quad \text{and} \quad \alpha_1 = \frac{3n-3}{2}.$$

Notice that $a_1 + (n+1) = \alpha_1$ and, by hypothesis, $\alpha_1 \geq n+1$. If $\alpha_1 = n+1$, we obtain $n = 5$, a contradiction. Otherwise, since $(n+1) \notin \lambda$, we obtain that λ is refinable. The claim (1) is then proved.

Let us now address the case (2). Similarly as before, we now have $a_2 = n-3, a_3 = n-2$ and $a_4 = n$ and so we determine

$$a_1 = \frac{n-3}{2} \quad \text{and} \quad \alpha_1 = \frac{3n-5}{2}.$$

Notice that from $\alpha_1 \geq n+1$ we obtain $n \geq 7$. However, assuming $n = 7$ leads to $a_1 + a_2 = 6 = n-1 \in \lambda$, a contradiction since λ is unrefinable. Let us now prove that the obtained partition

$$\lambda \stackrel{\text{def}}{=} \left(\dots, \widehat{\frac{n-3}{2}}, \dots, n-4, n-1, \frac{3n-5}{2}, 2n-4 \right)$$

is unrefinable by showing that each possible sum $a_i + a_j$, with $1 \leq i < j \leq 4$, is different from α_1 . Recall that by the classification of Theorem 4.1 we have already ruled out those partitions which contradict the unrefinability in $2n-4$. Since $n \geq 9$, we have that $a_1 + a_2 = (3n-9)/2 > n-1$. Consequently, every sum of missing parts is larger than $n-1 \in \lambda$. Moreover, $a_1 + a_2 \neq \alpha_1$, $a_1 + a_3 = (3n-7)/2 \neq \alpha_1$, $a_1 + a_4 = (3n-3)/2 > \alpha_1$, $a_2 + a_3 = 2n-5 > \alpha_1$ and therefore $a_2 + a_4, a_3 + a_4 > \alpha_1$. Therefore $\lambda \in \mathcal{C}_4$ and it is unique by construction, which proves the claim (2).

In the case of \mathcal{B}_4 , we find

$$a_1 = \frac{n-1}{2} \quad \text{and} \quad \alpha_1 = \frac{3n-7}{2}.$$

From $\alpha_1 \geq n+1$ we have $n \geq 9$, and assuming $n = 9$ contradicts again the unrefinability; therefore $n \geq 11$. With arguments similar to those of previous case the partition

$$\left(\dots, \widehat{\frac{n-1}{2}}, \dots, n-5, n-3, n, \frac{3n-7}{2}, 2n-4 \right)$$

is proved unrefinable and unique by construction, hence (3) is obtained.

Finally, considering the case of \mathcal{A}_4 , we obtain

$$a_1 = \frac{n+1}{2} \quad \text{and} \quad \alpha_1 = \frac{3n-9}{2}.$$

Now, $\alpha_1 \geq n+1$ implies $n \geq 11$ and $a_1 + a_2 = (3n-7)/2 > \alpha_1$. This proves that

$$\left(\dots, \widehat{\frac{n+1}{2}}, \dots, n-5, n-1, n, \frac{3n-9}{2}, 2n-4 \right) \in \mathcal{A}_4,$$

i.e. the claim (4). □

Proposition 5.5. *Let $n \geq 7$ be odd and $k \geq 0$. We have:*

- (1) if $n < 15$, then $\tilde{\mathcal{U}}_{T_n} \cap \mathcal{C}_5 = \emptyset$ and if $n \geq 15$, then $\#(\tilde{\mathcal{U}}_{T_{15+2k}} \cap \mathcal{C}_5) = \lfloor k/2 \rfloor + 1$,
- (2) if $n < 17$, then $\tilde{\mathcal{U}}_{T_n} \cap \mathcal{B}_5 = \emptyset$ and if $n \geq 17$, then $\#(\tilde{\mathcal{U}}_{17+2k} \cap \mathcal{B}_5) = \lfloor k/2 \rfloor + 1$,
- (3) if $n < 17$, then $\tilde{\mathcal{U}}_{T_n} \cap \mathcal{D}_5 = \emptyset$ and if $n \geq 17$, then $\#(\tilde{\mathcal{U}}_{17+2k} \cap \mathcal{D}_5) = \lfloor k/2 \rfloor + 1$,
- (4) if $n < 19$, then $\tilde{\mathcal{U}}_{T_n} \cap \mathcal{A}_5 = \emptyset$ and if $n \geq 19$, then $\#(\tilde{\mathcal{U}}_{19+2k} \cap \mathcal{A}_5) = \lfloor k/2 \rfloor + 1$.

Proof. Let us proceed as in the proof of Proposition 5.4. Let $\lambda \in \tilde{\mathcal{U}}_{T_n}$ be obtained by removing the integers a_1, a_2, \dots, a_5 and adding the replacements α_1, α_2 and $\alpha_3 = 2n-4$, which need to satisfy the following conditions:

- (i) $1 \leq a_1 < a_2 \leq n-5$,

- (ii) $n - 4 \leq a_3 < a_4 < a_5 \leq n$,
- (iii) $n + 1 \leq \alpha_2 = 2n - 4 - a_2 < \alpha_1 = 2n - 4 - a_1$,
- (iv) $a_1 + a_2 + a_3 + a_4 + a_5 = \alpha_1 + \alpha_2 + \alpha_3$.

First, let us address the case (1). We have $a_3 = n - 3, a_4 = n - 2$ and $a_5 = n$ and, from Eq. (iii) and Eq. (iv), a_1 and a_2 satisfy the condition

$$a_1 + a_2 = \frac{3n - 7}{2}. \quad (8)$$

We first consider the case when a_2 is maximal, i.e. $a_2 = n - 5$, in which we have $a_1 = (n + 3)/2$ and consequently $\alpha_1 = (3n - 11)/2$ and $\alpha_2 = n + 1$. Notice then that the condition of Eq. (8) can be met in other $\lfloor (a_2 - a_1 - 1)/2 \rfloor$ ways by taking the first two parts to be removed as $a_1 + i$ and $a_2 - i$, for $1 \leq i \leq \lfloor (a_2 - a_1 - 1)/2 \rfloor$. Now, from $a_1 < a_2$ we obtain $n \geq 15$. If $n = 15$, the partition

$$\left(\dots, \frac{\widehat{n+3}}{2}, \dots, \widehat{n-5}, n-4, n-1, n+1, \frac{3n-11}{2}, 2n-4 \right)$$

is unique by construction and is unrefinable since $a_1 + a_2 > \alpha_1$. In the other cases, which are

$$\left\lfloor \frac{a_2 - a_1 - 1}{2} \right\rfloor = \left\lfloor \frac{n - 15}{4} \right\rfloor, \quad (9)$$

we obtain an unrefinable partition since, letting $\alpha'_1 = 2n - 4 - (a_1 + i)$ and $\alpha'_2 = 2n - 4 - (a_2 - i)$, we have

$$(a_1 + i) + (a_2 - i) = a_1 + a_2 > \alpha_1 > \alpha'_1 > \alpha'_2.$$

The claim (1) is then obtained writing $n = 15 + 2k$ in Eq. (9).

The proofs for (2) and (4) are obtained in the same way. When $n = 17$, the partition

$$\left(\dots, \frac{\widehat{n+5}}{2}, \dots, \widehat{n-5}, n-3, n, n+1, \frac{3n-13}{2}, 2n-4 \right) \in \mathcal{B}_5$$

and is unique by construction, and when $n > 17$ it can be modified in $\lfloor (a_2 - a_1 - 1)/2 \rfloor = \lfloor (n - 17)/4 \rfloor$ ways as in the proof of (1). Analogously, when $n = 19$ the partition

$$\left(\dots, \frac{\widehat{n+7}}{2}, \dots, \widehat{n-5}, n-1, n, n+1, \frac{3n-15}{2}, 2n-4 \right) \in \mathcal{A}_5$$

and is unique by construction, and when $n > 19$ it can be modified in $\lfloor (n - 19)/4 \rfloor$ ways.

It remains to be proved the slightly different case (3). In this case, we have $a_1 + a_2 = (3n - 9)/2$ and, proceeding as above, from $a_2 = n - 5$ we obtain $a_1 = (n + 1)/2$ and $\alpha_1 = (3n - 9)/2$. This leads to the contradiction $a_1 + a_2 = \alpha_1$. The argument of (1) is here replicated starting from $a_2 = n - 6$. It is now easy to see that, when $n = 17$, the partition

$$\left(\dots, \frac{\widehat{n+3}}{2}, \dots, \widehat{n-6}, \dots, n-3, n+2, \frac{3n-11}{2}, 2n-4 \right),$$

unique by construction, is unrefinable. When $n > 17$, it can be modified in $\lfloor (n - 17)/4 \rfloor$ ways, which proves (3). \square

Proposition 5.6. *Let $n \geq 7$ be odd and $h \geq 6$. We have*

- (1) $\widetilde{\mathcal{U}}_{T_n} \cap \mathcal{D}_h \neq \emptyset$ if and only if $n \geq h^2 - h - 7$,
- (2) $\widetilde{\mathcal{U}}_{T_n} \cap \mathcal{C}_h \neq \emptyset$ if and only if $n \geq h^2 - h - 5$,
- (3) $\widetilde{\mathcal{U}}_{T_n} \cap \mathcal{B}_h \neq \emptyset$ if and only if $n \geq h^2 - h - 3$,

(4) $\tilde{\mathcal{U}}_{T_n} \cap \mathcal{A}_h \neq \emptyset$ if and only if $n \geq h^2 - h - 1$.

Proof. We proceed as in Proposition 5.4 and Proposition 5.5, assuming the conditions

- (i) $1 \leq a_1 < a_2 < \dots < a_{h-3} \leq n - 5$,
- (ii) $n - 4 \leq a_{h-2} < a_{h-1} < a_h \leq n$,
- (iii) $n + 1 \leq \alpha_{h-3} = 2n - 4 - a_{h-3} < \alpha_{h-4} = 2n - 4 - a_{h-4} < \dots < \alpha_1 = 2n - 4 - a_1$,
- (iv) $\sum a_i = \sum \alpha_i$.

If $\lambda \in \tilde{\mathcal{U}}_{T_n} \cap \mathcal{D}_h$, then $a_{h-2} + a_{h-1} + a_h = 3n - 3$ and therefore, from Eq. (iii) and Eq. (iv),

$$a_1 + a_2 + \dots + a_{h-3} = \frac{(h-2)(2n-4) - (3n-3)}{2} = \frac{(2h-7)n + 11 - 4h}{2}.$$

Let us now assume that $a_{h-3} = n - 5, a_{h-4} = n - 6, \dots, a_2 = n - h$, i.e. let us maximize the sum $a_2 + \dots + a_{h-3}$. We obtain

$$a_2 + \dots + a_{h-3} = (h-4)n - \sum_{i=5}^h i = (h-4)n - \frac{h(h+1)}{2} + 10,$$

from which we can calculate

$$a_1 = \frac{n + h^2 - 3h - 9}{2}.$$

Imposing $a_1 < a_2$ we obtain $n > h^2 - h - 9$. In this setting, we have

$$\alpha_1 = \frac{3n - h^2 + 3h + 1}{2} \quad \text{and} \quad a_1 + a_2 = \frac{3n + h^2 - 5h - 9}{2}.$$

Notice that $a_1 + a_2 > \alpha_1$ is satisfied for $h \geq 6$, hence the provided construction leads to a partition λ which belongs to \mathcal{D}_h if and only if $n \geq h^2 - h - 7$, i.e. (1).

The cases (2), (3) and (4) we proceed analogously, maximizing $a_2 + a_3 + \dots + a_{h-3}$, provided that a_{h-2}, a_{h-1}, a_h are modified accordingly. In particular, when considering \mathcal{C}_h we have

$$a_1 = \frac{n + h^2 - 3h - 7}{2} \quad \text{and} \quad \alpha_1 = \frac{3n - h^2 + 3h - 1}{2}.$$

From $a_1 < a_2$ we have $n \geq h^2 - h - 5$ and from

$$\alpha_1 < a_1 + a_2 = \frac{3n + h^2 - 5h - 7}{2}$$

we obtain $\lambda \in \mathcal{C}_h$, i.e. the claim (2) follows.

In the case of \mathcal{B}_h we have

$$a_1 = \frac{n + h^2 - 3h - 5}{2} \quad \text{and} \quad \alpha_1 = \frac{3n - h^2 + 3h - 3}{2}.$$

From $a_1 < a_2$ we have $n \geq h^2 - h - 3$ and from

$$\alpha_1 < a_1 + a_2 = \frac{3n + h^2 - 5h - 5}{2}$$

we obtain $\lambda \in \mathcal{B}_h$, i.e. the claim (3) is proved.

Finally, assuming the conditions of \mathcal{A}_h we have

$$a_1 = \frac{n + h^2 - 3h - 3}{2} \quad \text{and} \quad \alpha_1 = \frac{3n - h^2 + 3h - 5}{2}.$$

From $a_1 < a_2$ we have $n \geq h^2 - h - 1$ and from

$$\alpha_1 < a_1 + a_2 = \frac{3n + h^2 - 5h - 3}{2}$$

we obtain $\lambda \in \mathcal{A}_h$, from which the desired result (4) follows. \square

By interchanging the role of n and h in the statements of Proposition 5.6, we obtain the following description of the set of maximal unrefinable partitions of triangular numbers of an odd number, where we can read the upper bound for h in each different class.

Corollary 5.7. *Let $n \geq 7$ be odd. Then*

$$\tilde{\mathcal{U}}_{T_n} = \{\widetilde{\pi}_n\} \dot{\cup} \left(\bigcup_{h=5}^{\lfloor \frac{1+\sqrt{29+4n}}{2} \rfloor} \mathcal{D}_h \dot{\cup} \bigcup_{h=4}^{\lfloor \frac{1+\sqrt{21+4n}}{2} \rfloor} \mathcal{C}_h \dot{\cup} \bigcup_{h=4}^{\lfloor \frac{1+\sqrt{13+4n}}{2} \rfloor} \mathcal{B}_h \dot{\cup} \bigcup_{h=4}^{\lfloor \frac{1+\sqrt{5+4n}}{2} \rfloor} \mathcal{A}_h \right) \cap \tilde{\mathcal{U}}_{T_n}.$$

Remark 4. In the proof of Proposition 5.6 we have exhibited an example of unrefinable partition for each class, constructed by maximizing $a_2 + a_3 + \dots + a_{h-3}$ and consequently by determining a_1 . The unrefinability of the obtained partition is then granted from the fact that $a_1 + a_2 > \alpha_1$. Notice that, each other partition λ' of the same class is determined by the removed parts $a'_1, a'_2, \dots, a'_{h-3}$ such that $a'_1 = a_1 + i$ and $a'_s = a_s - i_{s-1}$ for $s > 1$ and $i_s \geq 0$, where $i = \sum_{s=1}^{h-4} i_s$, provided that $a'_i < a'_s$ for $i < s$. The unrefinability of λ' is then easily proved, since

$$a'_1 + a'_2 = a_1 + i + a_2 - i_i \geq a_1 + a_2 \geq \alpha_1 \geq \alpha'_1.$$

Example 5.8. Let $n = 49$. For the bound in the previous corollary, when considering partitions of class \mathcal{D} we have $5 \leq h \leq (1 + \sqrt{29 + 4n})/2 = 8$. Let us fix $h = 7$ and construct all the partitions in $\tilde{\mathcal{U}}_{T_{49}} \cap \mathcal{D}_7$. We recall that, for Theorem 4.1, a partition of class \mathcal{D}_7 is given when $a_1, a_2, \dots, a_{h-3} = a_4$ are specified. Therefore, for sake of simplicity, we denote the partitions just by listing the removed parts (a_1, a_2, a_3, a_4) . Let us start, as in Proposition 5.6, from the partition

$$\left(\frac{n + h^2 - 3h - 9}{2}, n - 7, n - 6, n - 5 \right) = (34, 42, 43, 44).$$

All the remaining partitions in \mathcal{D}_7 , obtained as in Remark 4, are:

$$\begin{aligned} &(35, 41, 43, 44) \quad (36, 40, 43, 44) \\ &(37, 39, 43, 44) \quad (36, 41, 42, 44) \\ &(37, 40, 42, 44) \quad (38, 39, 42, 44) \\ &(38, 40, 41, 44) \quad (37, 41, 42, 43) \\ &(38, 40, 42, 43) \quad (39, 40, 41, 43) \end{aligned}$$

The partitions in other classes are obtained analogously.

We have already highlighted in Example 5.2 and in Example 5.3 what $\min_{\lambda \in \tilde{\mathcal{U}}_{T_n}} \text{mex}(\lambda)$ looks like. The intuition can now be easily proved as a consequence of the previous propositions.

Corollary 5.9. *Let $n \geq 7$ be odd. For each $\lambda \in \tilde{\mathcal{U}}_{T_n}$ we have*

$$\text{mex}(\lambda) = \mu_1 \geq \frac{(n-3)}{2}.$$

Proof. Notice that $\mu_1 = a_1$. The claim is trivial if $\lambda = \widetilde{\pi}_n$. Otherwise it follows from Proposition 5.4, 5.5 and 5.6, recalling that a_1 was calculated in order to be minimal, since $a_2 + a_3 + \dots + a_{h-3}$ was maximized. The results are summarized in Table 3, where is not hard to check that $(n-3)/2$ is the smaller value that a_1 can assume. \square

class	a_1
\mathcal{C}_4	$(n-3)/2$
\mathcal{B}_4	$(n-1)/2$
\mathcal{A}_4	$(n+1)/2$
\mathcal{D}_5	$(n+3)/2$
\mathcal{C}_5	$(n+3)/2$
\mathcal{B}_5	$(n+5)/2$
\mathcal{A}_5	$(n+7)/2$
\mathcal{D}_h	$(n+h^2-3h-9)/2$
\mathcal{C}_h	$(n+h^2-3h-7)/2$
\mathcal{B}_h	$(n+h^2-3h-5)/2$
\mathcal{A}_h	$(n+h^2-3h-3)/2$

TABLE 3. Values of a_1 in the construction of Proposition 5.4, 5.5 and 5.6, for $h = 4$, $h = 5$ and $h \geq 6$.

5.2. The bijection. In this conclusive section we prove the main contribution of this work, i.e. we show that, when n is odd, the number of maximal unrefinable partitions of T_n equals the number of partitions of $\lceil n/2 \rceil$ into distinct parts by means of a bijective proof. Notice that, by the anti-symmetric property (Remark 3) and by the bound on the minimal excludant (Corollary 5.9), a partition in $\widetilde{\mathcal{U}}_{T_n}$ is determined by at most

$$(n-2) - \frac{n-3}{2} = \frac{n+1}{2} - 1 = \left\lfloor \frac{n}{2} \right\rfloor - 1$$

parts, and so is a partition of $\lceil n/2 \rceil$ into distinct parts. The following theorem is used to establish a bijection between $\widetilde{\mathcal{U}}_{T_{2k-1}}$ and \mathbb{D}_k .

Theorem 5.10. *Let a_1, a_2, \dots, a_u be the missing parts smaller or equal than $n-3$ of an unrefinable partition $\lambda \neq \widetilde{\pi}_n$ of T_n , for some odd integer $n \geq 7$. Then n , λ and its class are uniquely determined.*

Proof. Let us start by proving that n can be obtained by the knowledge of a_1, a_2, \dots, a_u . In particular, let us prove that

$$n = \frac{2 \sum_{i=1}^u a_i + 1 + 4u}{2u-1} \quad (10)$$

by distinguishing the four possible classes. Let us first assume $\lambda \in \mathcal{D}_h$ for some $h \geq 5$. Recalling that $a_u \leq n-3$ and, by definition of \mathcal{D}_h and by Remark 3, since $a_{h-1} = n-1$ and $a_h = n$, we have $a_u \notin \{n-3, n-4\}$. Therefore $a_u \leq n-5$, and so $u = h-3$. Recalling that the following conditions hold

- (i) $1 \leq a_1 < a_2 < \dots < a_{h-3} \leq n-5$,
- (ii) $n-4 \leq a_{h-2} < a_{h-1} < a_h \leq n$,
- (iii) $n < \alpha_{h-3} = 2n-4-a_{h-3} < \alpha_{h-4} = 2n-4-a_{h-4} < \dots < \alpha_1 = 2n-4-a_1$,
- (iv) $\sum a_i = \sum \alpha_i$,

we obtain

$$\sum_{i=1}^u a_i = \frac{(u+1)(2n-4) - (3n-3)}{2},$$

from which we determine n as claimed.

Let us consider the class \mathcal{C}_h . In this case, reasoning as above, we have $a_u = n - 3$ and $a_{u-1} \leq n - 5$, which means $u - 1 = h - 3$. Therefore

$$\sum_{i=1}^u a_i - (n - 3) = \sum_{i=1}^{u-1} a_i = \frac{u(2n - 4) - (3n - 5)}{2},$$

from which we obtain again Eq. (10).

When $\lambda \in \mathcal{B}_h$, we have $a_u = n - 4$, which means $h = u + 2$, so

$$\sum_{i=1}^u a_i - (n - 4) = \sum_{i=1}^{u-1} a_i = \frac{u(2n - 4) - (3n - 7)}{2},$$

from which the same n is determined.

In conclusion, if $\lambda \in \mathcal{A}_h$, we have $a_u = n - 3$ and $a_{u-1} = n - 4$, so $h = u + 1$ and Eq. (10) is satisfied since

$$\sum_{i=1}^u a_i - (n - 4) - (n - 3) = \sum_{i=1}^{u-2} a_i = \frac{(u - 1)(2n - 4) - (3n - 9)}{2}.$$

Now that n is determined from a_1, a_2, \dots, a_u , the class of the partition can be recognized by looking at a_{u-1} and a_u . In particular

- $a_u < n - 4 \iff \lambda \in \mathcal{D}_{u+3}$,
- $a_u = n - 3$ and $a_{u-1} < n - 4 \iff \lambda \in \mathcal{C}_{u+2}$,
- $a_u = n - 4 \iff \lambda \in \mathcal{B}_{u+2}$,
- $a_u = n - 3$ and $a_{u-1} = n - 4 \iff \lambda \in \mathcal{A}_{u+1}$.

To conclude, we determine the partition by using the anti-symmetric property (cf. Remark 3). \square

We are now ready to prove our last result. Denoting by \mathbb{D} the set of all the partitions into distinct parts, let us define the following subsets of \mathbb{D} :

$$\begin{aligned} \mathcal{A}_t^* &\stackrel{\text{def}}{=} \{\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t) \mid \lambda \in \mathbb{D}, \lambda_1 = 1, \lambda_2 = 2, t \geq 3\}, \\ \mathcal{B}_t^* &\stackrel{\text{def}}{=} \{\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t) \mid \lambda \in \mathbb{D}, \lambda_1 = 2, t \geq 2\}, \\ \mathcal{C}_t^* &\stackrel{\text{def}}{=} \{\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t) \mid \lambda \in \mathbb{D}, \lambda_1 = 1, \lambda_2 > 2, t \geq 2\}, \\ \mathcal{D}_t^* &\stackrel{\text{def}}{=} \{\lambda = (\lambda_1, \lambda_2, \dots, \lambda_t) \mid \lambda \in \mathbb{D}, \lambda_1 > 3, t \geq 2\}. \end{aligned}$$

It is not hard to notice that

$$\mathbb{D} = \mathcal{A}^* \dot{\cup} \mathcal{B}^* \dot{\cup} \mathcal{C}^* \dot{\cup} \mathcal{D}^*,$$

where

$$\mathcal{A}^* \stackrel{\text{def}}{=} \bigcup_{t \geq 3} \mathcal{A}_t^*, \quad \mathcal{B}^* \stackrel{\text{def}}{=} \bigcup_{t \geq 2} \mathcal{B}_t^*, \quad \mathcal{C}^* \stackrel{\text{def}}{=} \bigcup_{t \geq 2} \mathcal{C}_t^*, \quad \mathcal{D}^* \stackrel{\text{def}}{=} \bigcup_{t \geq 2} \mathcal{D}_t^*.$$

Let us conclude the paper proving a bijection between $\widetilde{\mathbb{U}}_{T_{2k-1}}$ and \mathbb{D}_k .

Theorem 5.11. *Let $k \in \mathbb{N}$, $k \geq 7$, $n = 2k - 1$ and $N = T_n$. Let $\sigma: \widetilde{\mathbb{U}}_{T_n} \rightarrow \mathbb{D}_k$ be such that*

$$\lambda \mapsto \begin{cases} (3, k - 3) & \lambda = \widetilde{\pi}_n \\ (n - 2 - a_u, \dots, n - 2 - a_2, n - 2 - a_1) & \lambda \neq \widetilde{\pi}_n \end{cases},$$

where, if $\lambda \neq \widetilde{\pi}_n$, then (a_1, a_2, \dots, a_u) are the missing parts of $\mathcal{M}_\lambda \cap \{1, 2, \dots, n - 3\}$ as in Theorem 5.10. Then σ is bijective, therefore $\#\widetilde{\mathbb{U}}_{T_{2k-1}} = \#\mathbb{D}_k$.

Proof. For sake of brevity and by virtue of Theorem 5.10, we will denote each $\lambda \in \widetilde{\mathcal{U}}_{T_n} \setminus \{\widetilde{\pi}_n\}$ by listing its missing parts a_1, a_2, \dots, a_u in $\mathcal{M}_\lambda \cap \{1, 2, \dots, n-3\}$. We prove that σ is bijective by proving explicitly that partitions of $\mathcal{A} \cap \widetilde{\mathcal{U}}_{T_{2k-1}}$ are in one-to-one correspondence with those of $\mathcal{A}^* \cap \mathbb{D}_k$ and that the same holds respectively for \mathcal{B} and \mathcal{B}^* , \mathcal{C} and \mathcal{C}^* , and $\mathcal{D} \cup \{\widetilde{\pi}_n\}$ and \mathcal{D}^* .

Let us start by proving that σ is well defined, i.e. for each $\lambda \in \widetilde{\mathcal{U}}_{T_n}$ we have that $\sigma(\lambda)$ is a partition of k into distinct parts. If $\lambda = \widetilde{\pi}_n$ there is nothing to prove, otherwise, since the missing parts of λ are distinct, so are the parts $n-2-a_u < \dots < n-2-a_2 < n-2-a_1$ of $\sigma(\lambda)$. We now prove that the sum of the parts of $\sigma(\lambda)$ is k in each possible case, making extensively use of Proposition 5.4, Proposition 5.5 and Proposition 5.6 without further mention.

If $\lambda \in \mathcal{A}_4$, then $\lambda = ((n+1)/2, n-4, n-3)$ and so

$$\sigma(\lambda) = \left(1, 2, n-2 - \frac{n+1}{2}\right) = \left(1, 2, \frac{n-5}{2}\right) = \left(1, 2, \frac{n+1}{2} - 3\right) = (1, 2, k-3) \in \mathbb{D}_k.$$

Notice that, in particular, $\sigma(\lambda) \in \mathbb{D}_k \cap \mathcal{A}_3^*$. Similarly, if $\lambda \in \mathcal{B}_4$, then $\lambda = ((n-1)/2, n-4)$ and $\sigma(\lambda) = (2, (n+1)/2 - 2) = (2, k-2) \in \mathbb{D}_k \cap \mathcal{B}_2^*$. If $\lambda \in \mathcal{C}_4$ we have $\lambda = ((n-3)/2, n-3)$, so $\sigma(\lambda) = (1, (n+1)/2 - 1) = (1, k-1) \in \mathbb{D}_k \cap \mathcal{C}_2^*$.

If $\lambda \in \mathcal{A}_5$, then $\lambda = ((n+7)/2 + i, n-5-i, n-4, n-3)$, for $0 \leq i \leq \lfloor (n-19)/4 \rfloor$, and

$$\sigma(\lambda) = \left(1, 2, 3+i, \frac{n-11}{2} - i\right) = \left(1, 2, 3+i, \frac{n+1}{2} - 6 - i\right) = (1, 2, 3+i, k-6-i) \in \mathbb{D}_k,$$

for $0 \leq i \leq \lfloor (n-19)/4 \rfloor$. In particular, $\sigma(\lambda) \in \mathbb{D}_k \cap \mathcal{A}_4^*$. Similarly, if $\lambda \in \mathcal{B}_5$, then $\lambda = ((n+5)/2 + i, n-5-i, n-4)$ and

$$\sigma(\lambda) = \left(2, 3+i, \frac{n+1}{2} - 5 - i\right) = (2, 3+i, k-5-i) \in \mathbb{D}_k \cap \mathcal{B}_3^*,$$

for $0 \leq i \leq \lfloor (n-17)/4 \rfloor$. If $\lambda \in \mathcal{C}_5$, we have $\lambda = ((n+3)/2 + i, n-5-i, n-3)$ and so

$$\sigma(\lambda) = \left(1, 3+i, \frac{n+1}{2} - 4 - i\right) = (1, 3+i, k-4-i) \in \mathbb{D}_k \cap \mathcal{C}_3^*,$$

for $0 \leq i \leq \lfloor (n-15)/4 \rfloor$. In the case when $\lambda = ((n+3)/2 + i, n-6-i) \in \mathcal{D}_5$, we have

$$\sigma(\lambda) = \left(4+i, \frac{n+1}{2} - 4 - i\right) = (4+i, k-4-i) \in \mathbb{D}_k \cap \mathcal{D}_2^*,$$

for $0 \leq i \leq \lfloor (n-17)/4 \rfloor$.

Let us now consider $\lambda \in \mathcal{D}_h$ for $h \geq 6$. In this case

$$\lambda = ((n+h^2-3h-9)/2 + i, n-h-i_1, \dots, n-5-i_{h-4}),$$

where $i = \sum_{s=1}^{h-4} i_s$, and so

$$\begin{aligned} \sigma(\lambda) &= \left(3+i_{h-4}, \dots, h-2+i_1, \frac{n-h^2+3h+5}{2} - i\right) \\ &= \left(3+i_{h-4}, \dots, h-2+i_1, k+2 - \frac{h^2-3h}{2} - i\right). \end{aligned}$$

Notice that

$$\begin{aligned}
 & (3 + i_{h-4}) + (4 + i_{h-5}) + \cdots + (h - 2 + i_1) + \\
 & \quad + \left(k + 2 - \frac{h^2 - 3h}{2} - i \right) = \\
 & \quad \sum_{j=3}^{h-2} j + \sum_{s=1}^{h-4} i_s + k + 2 - \frac{h^2 - 3h}{2} - i = \\
 & \quad \frac{(h-2)(h-1)}{2} - 3 + k + 2 - \frac{h^2 - 3h}{2} = \\
 & \quad \frac{h^2 - 3h}{2} - 2 + k + 2 - \frac{h^2 - 3h}{2} = k,
 \end{aligned}$$

and so $\sigma(\lambda) \in \mathbb{D}_k \cap \mathcal{D}_{h-3}^*$. Similarly, if $\lambda \in \mathcal{C}_h$, then

$$\lambda = \left(\frac{n + h^2 - 3h - 7}{2} + i, n - h - i_1, \dots, n - 5 - i_{h-4}, n - 3 \right)$$

and

$$\sigma(\lambda) = \left(1, 3 + i_{h-4}, \dots, h - 2 + i_1, k + 1 - \frac{h^2 - 3h}{2} - i \right) \in \mathbb{D}_k \cap \mathcal{C}_{h-2}^*$$

since the sum of the first $h - 3$ terms is $(h^2 - 3h)/2 - 1 + \sum_{s=1}^{h-4} i_s$. If $\lambda \in \mathcal{B}_h$, we have

$$\lambda = \left(\frac{n + h^2 - 3h - 5}{2} + i, n - h - i_1, \dots, n - 5 - i_{h-4}, n - 4 \right)$$

and so

$$\sigma(\lambda) = \left(2, 3 + i_{h-4}, \dots, h - 2 + i_1, k - \frac{h^2 - 3h}{2} - i \right) \in \mathbb{D}_k \cap \mathcal{B}_{h-2}^*,$$

since the sum of the first $h - 3$ terms is $(h^2 - 3h)/2 + \sum_{s=1}^{h-4} i_s$. Finally, in the case when

$$\lambda = \left(\frac{n + h^2 - 3h - 3}{2} + i, n - h - i_1, \dots, n - 5 - i_{h-4}, n - 4, n - 3 \right) \in \mathcal{A}_h, \quad (11)$$

we obtain

$$\sigma(\lambda) = \left(1, 2, 3 + i_{h-4}, \dots, h - 2 + i_1, k - 1 - \frac{h^2 - 3h}{2} - i \right) \in \mathbb{D}_k \cap \mathcal{A}_{h-1}^*, \quad (12)$$

noticing that the sum of the first $h - 2$ terms is $(h^2 - 3h)/2 + 1 + \sum_{s=1}^{h-4} i_s$.

We proved that σ is well defined. Notice also that σ is trivially injective. Therefore it remains to prove that σ is surjective. In particular, it suffices to check that for each partition $\lambda^* \in (\mathcal{A}^* \dot{\cup} \mathcal{B}^* \dot{\cup} \mathcal{C}^* \dot{\cup} \mathcal{D}^*) \cap \mathbb{D}_k$, $\lambda^* \neq (3, k - 3)$, there exists $\lambda \in (\mathcal{A} \dot{\cup} \mathcal{B} \dot{\cup} \mathcal{C} \dot{\cup} \mathcal{D}) \cap \widetilde{\mathbb{U}}_{T_{2k-1}}$ such that $\sigma(\lambda) = \lambda^*$, since $\sigma(\widetilde{\pi}_n) = (3, k - 3)$ by definition. Given $\lambda^* = (\lambda_1^*, \lambda_2^*, \dots, \lambda_t^*) \in \mathbb{D}_k$, by definition of σ we have that the partition λ denoted by its missing parts $(n - 2 - \lambda_t^*, \dots, n - 2 - \lambda_2^*, n - 2 - \lambda_1^*)$ is such that $\sigma(\lambda) = \lambda^*$. It remains to prove that such λ is a maximal unrefinable partition of n . The full details of the proof are here omitted since they can be obtained by arguments very similar to those used for proving that σ is well defined. As an example, let us consider the case when $\lambda^* \in \mathcal{A}_t^* \cap \mathbb{D}_k$, for $t \geq 5$, and let us prove that λ^* is the image of an unrefinable partition λ of class \mathcal{A} . Since λ^* is a partition of k into t distinct parts and containing 1 and 2 by definition we can write

$$\lambda^* = \left(1, 2, 3 + i_1, \dots, t - 1 + i_{t-3}, k - \sum_{s=1}^{t-1} \lambda_s^* \right), \quad (13)$$

x										$n-2$	n										λ_t
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22
				6	5	4	3	2	1												
•	•	•	•	•	•	•	•	•	•		•	•		•							•
•	•	•	•	•	•	•	•	•	•			•			•						•
•	•	•	•	•	•	•	•	•	•		•					•					•

 TABLE 4. The bijection σ shown on the partitions of $\lambda \in \widetilde{\mathcal{U}}_{T_{13}}$, $\lambda \neq \widetilde{\pi}_{13}$.

where

$$\sum_{s=1}^{t-1} \lambda_s^* = \frac{(t-1)t}{2} + \sum_{s=1}^{t-3} i_s$$

for some $i_1, i_2, \dots, i_{t-3} \geq 0$ (cf. also Eq. (12)). We can now substitute $t-1$ to $h-2$ and $(n+1)/2$ to k in Eq. (13). Applying the correspondence $\lambda_i \leftrightarrow n-2 - \lambda_{t-i+1}^*$ and denoting the obtained partition λ by listing its missing parts, we obtain

$$\lambda = \left(\frac{n+h^2-3h-3}{2} + i, n-h-i_{h-4}, \dots, n-5-i_1, n-4, n-3 \right)$$

as in Eq. (11). This proves that $\lambda \in \widetilde{\mathcal{U}}_{T_{2k-1}} \cap \mathcal{A}_{t+1}$ is unrefinable (cf. Proposition 5.6 and Remark 4) and such that $\sigma(\lambda) = \lambda^*$. The remaining cases are similar. \square

Remark 5. The bijection σ is not well defined when $k < 7$. However, it can be easily shown that the result of Theorem 5.11 is still valid when $k = 4$ and $k = 5$, where we have $\#\widetilde{\mathcal{U}}_{T_7} = \#\mathbb{D}_4 = 1$ and $\#\widetilde{\mathcal{U}}_{T_5} = \#\mathbb{D}_5 = 2$, respectively. The claim is false instead in the case $k = 6$, where we have $\#\widetilde{\mathcal{U}}_{T_{11}} = 4$ and $\#\mathbb{D}_6 = 3$.

In the proof of Theorem 5.11 we showed that σ is a bijection from $\widetilde{\mathcal{U}}_{T_{2k-1}}$ to \mathbb{D}_k . More than this, we also proved that σ is bijective when restricted to each class.

Corollary 5.12. *The function σ of Theorem 5.11 sends in a bijective way*

- (i) $\mathcal{A}_h \cap \widetilde{\mathcal{U}}_{T_{2k-1}}$ to $\mathcal{A}_{h-1}^* \cap \mathbb{D}_k$,
- (ii) $\mathcal{B}_h \cap \widetilde{\mathcal{U}}_{T_{2k-1}}$ to $\mathcal{B}_{h-2}^* \cap \mathbb{D}_k$,
- (iii) $\mathcal{C}_h \cap \widetilde{\mathcal{U}}_{T_{2k-1}}$ to $\mathcal{C}_{h-2}^* \cap \mathbb{D}_k$,
- (iv) $\mathcal{D}_h \cap \widetilde{\mathcal{U}}_{T_{2k-1}}$ to $(\mathcal{D}_{h-3}^* \setminus \{(3, k-3)\}) \cap \mathbb{D}_k$.

Example 5.13. Coming back to the case of Example 5.2, we represent in Tab. 4 the bijection σ between maximal unrefinable partitions of 13 obtained in the case $h = j-2$ (hence those different from $\widetilde{\pi}_{13}$), represented by black dots, and the partitions of 7 into distinct parts, represented by blue dots. Notice that the partition $(3, 4)$ is not displayed since it corresponds to $\widetilde{\pi}_{13}$. Here x corresponds to

$$x \stackrel{\text{def}}{=} \min_{\lambda \in \widetilde{\mathcal{U}}_{T_{13}}} \text{mex}(\lambda).$$

Equivalently, by the anti-symmetric property, partitions of 7 into distinct parts can be read looking at the black dots in the right side of the table.

Remark 6. Another combinatorial equality can be derived from the provided construction for $\widetilde{\mathcal{U}}_N$. Indeed, assuming $n = 2k-1$ for $k \geq 7$, $h \geq 6$, and reasoning as in Example 5.8,

class	$f(n, h)$	$g(n, h)$
\mathcal{A}_h	$\frac{-h^3+6h^2+(n-8)h-4n+2}{2}$	$\frac{n-h^2+3h-5}{2}$
\mathcal{B}_h	$\frac{-h^3+6h^2+(n-6)h-4n-6}{2}$	$\frac{n-h^2+3h-3}{2}$
\mathcal{C}_h	$\frac{-h^3+6h^2+(n-4)h-4n-14}{2}$	$\frac{n-h^2+3h-1}{2}$
\mathcal{D}_h	$\frac{-h^3+6h^2+(n-2)h-4n-22}{2}$	$\frac{n-h^2+3h+1}{2}$

 TABLE 5. The values of $f(n, h)$ and $g(n, h)$ for each class.

it can be easily shown that $\#(\tilde{\mathcal{U}}_{T_n} \cap \mathcal{D}_h)$ equals the number of partitions in $h-3$ parts of $f(n, h)$ in which each part is smaller or equal than $g(n, h)$, where

$$f(n, h) = \frac{-h^3 + 6h^2 + (n-2)h - 4n - 22}{2} \quad \text{and} \quad g(n, h) = \frac{n - h^2 + 3h + 1}{2}.$$

The proof is obtained from Proposition 5.6, considering the bijection

$$a_i \leftrightarrow a_i - a_1 + 1. \quad (14)$$

In Tab. 6 the result is summarized for each class. Notice that, using the bijection of Eq. (14) on the partitions shown in Example 5.8, one can recover the eleven partitions of 31 in 4 parts, where each part is not larger than 11.

REFERENCES

- [ACGS19] R. Aragona, R. Civino, N. Gavioli, and C. M. Scoppola, *Regular subgroups with large intersection*, Ann. Mat. Pura Appl. (4) **198** (2019), no. 6, 2043–2057.
- [ACGS21a] ———, *Rigid commutators and a normalizer chain*, Monatsh. Math. **196** (2021), no. 3, 431–455.
- [ACGS21b] ———, *Unrefinable partitions into distinct parts in a normalizer chain*, arXiv preprint arXiv:2107.04666 (2021).
- [AN19] G. E. Andrews and D. Newman, *Partitions and the minimal excludant*, Ann. Comb. **23** (2019), no. 2, 249–254.
- [And76] G. E. Andrews, *The theory of partitions*, Encyclopedia of Mathematics and its Applications, Vol. 2, Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1976.
- [FP15] A. S. Fraenkel and U. Peled, *Harnessing the unwieldy MEX function*, Games of no chance 4, Math. Sci. Res. Inst. Publ., vol. 63, Cambridge Univ. Press, New York, 2015, pp. 77–94.
- [HSS22] B. Hopkins, J. A. Sellers, and D. Stanton, *Dyson’s crank and the mex of integer partitions*, J. Combin. Theory Ser. A **185** (2022), Paper No. 105523, 10.
- [OEI] *The On-Line Encyclopedia of Integer Sequences*, Published electronically at <https://oeis.org>, Accessed: 2021-11-01.